

# Vermicomposting: a technology for vermiremediation of heavy metals from sewage sludge and animal dung

Deepak Kumar Bhartiya<sup>1</sup> , Gorakh Nath<sup>2</sup> , Keshav Singh<sup>3,\*</sup> 

<sup>1</sup>Department of Zoology, Government Degree College, Dhadha Bujurg, Hata, Kushinagar, U.P., India.

<sup>2</sup>Department of Zoology, S.V.M. Mahila, P.G. College, Gorakhpur, U.P., India.

<sup>3</sup>Vermibiotechnology Laboratory, Department of Zoology, D.D.U. Gorakhpur University, Gorakhpur, U.P., India.

\*Corresponding author: [keshav26singh@rediffmail.com](mailto:keshav26singh@rediffmail.com)

## Original Research

## Abstract:

Received:

12 Apr 2023

Revised:

11 Aug 2023

Accepted:

13 Dec 2023

Published online:

20 May 2024

© The Author(s) 2024

**Purpose:** The present study to investigate the earthworm *Eisenia fetida* was a chief organism for accumulation of heavy metals (Pb, Cr, and Cd) in their body tissue from sewage sludge with different combinations of animal dung and converted into rich organic vermicompost which plays a beneficial role for environment, animals and human health.

**Method:** Animal dung and sewage sludge were collected from Gorakhpur city and earthworm *E. fetida* an epigeic species collected from Vermibiotechnology Laboratory, Department of Zoology, D. D. U. Gorakhpur University, Gorakhpur. Analysis of heavy metals in different animal dung, sewage sludge, final vermicompost, and earthworm bodies were measured by Shimadzu AA- 7000 atomic absorption spectrometer.

**Results:** There was a significant ( $P < 0.05$ ) reduction of Pb, Cr, and Cd concentration observed in different animal dung such as cow, buffalo and goat dung with sewage sludge in ratios of 1:1, 1:2, and 1:3 before and after vermicomposting. Cr was significantly decreased in cow dung (CD) and goat dung (GD) and Cd in cow dung (CD) and cow dung with sewage sludge (CD+ SS, 1:1) the at BDL level whereas the concentration of Pb significantly decreased by 76.470% in the combination of GD+ SS (1:1). The Cr and Cd were significantly increased in the body of *E. fetida* 3.384% and 11.801%, respectively, in the combination of CD+ SS (1:3) whereas Pb significantly increase by 23.018% in the combination of BD+SS (1:3).

**Conclusion:** *E. fetida* is a suitable species for the accumulation of heavy metals such as Pb, Cr, and Cd in different vermibeds during vermicomposting and plays an important role in the remediation of heavy metals from animal dung with sewage sludge. Vermibiotechnology is a useful technique for the management of heavy metals.

**Keywords:** Vermiremediation; *Eisenia fetida*; Sewage sludge; Heavy metals; Vermicomposting

## 1. Introduction

The production of sewage sludge is parallel to the increasing rate of population in India. It is a serious problem for society and it causes environmental hazards (Kaviraj and Sharma, 2003; Suthar, 2006; Yadav and Garg, 2011; Jatav et al., 2021). Cow, buffalo, sheep, goat, and horse excreta cause odor and pollution problems (Gupta, 2005; Garg et al., 2006; Nath et al., 2009). Animal dung and sewage sludge contain heavy metals that cause ill effects on soil as well as human health (Prajapati and Singh, 2013; Rusanescu et al., 2022). Heavy metals are generally referred to as those metals which possess a specific density

of more than 5 g/cm<sup>3</sup> and adversely affect the environment and living organisms (Jarup, 2003). The safe value of Lead is 0.03 mg/kg, Chromium is 0.01 mg/kg and according to the FAO/WHO, the recommended tolerable daily Cd uptake is 0.01 to 0.035 mg/d (WHO, 1985, 1993). The toxic metals, persistent organic compounds, pathogenic bacteria, and parasite eggs, could pollute the soil and groundwater (Dume et al., 2022). High exposure to lead causes severe damage to the brain and kidneys, anemia, increased blood pressure, miscarriage in pregnant women, and reduced fertility in males (Cleveland et al., 2008). Ahmed et al. (2016) reported that chromium eventually accumulates in crops from contaminated soils and imparts health risks to humans

and animals, through the food chain. Cadmium is primarily toxic to the kidney and it causes damage to the proximal tubules and failure of the tubular reabsorption leading to renal dysfunction (Lee et al., 2006).

Vermicomposting of sewage sludge with animal dung by *Eisenia fetida* is an ecological method of sewage sludge management (Spurgeon and Hopkin, 1999; Iwai et al., 2013). The worms break down these organic wastes and modify them into nutrient-rich organic manure i.e. vermicompost. *E. fetida* is often used in vermicomposting as it reduces the C/N ratio, the pathogenic load of various wastes but increases the total Kjeldahl nitrogen (TKN) and total phosphorus (TP), hence obtaining good quality fertilizer for soil (Li et al., 2011; Iwai et al., 2013; Panday et al., 2014). Earthworm *E. fetida* biochemically degrades sewage sludge and homogenizes the material through the muscular actions of the intestine and releases nutrient-rich vermicompost for plants (Yadav et al., 2012; Purdea et al., 2019) by accumulating heavy metals in their body during the vermicomposting (Suleiman et al., 2017; Singh and Bhartiya, 2021). The aim of the present study was to investigate the accumulation of heavy metals in earthworm *E. fetida* in vermicomposting of animal dung with sewage sludge and converted rich organic vermicompost which is ecologically sound and culturally most acceptable for agriculture.

## 2. Materials and methods

### Collection of animal dung, earthworm, and sewage sludge (SS)-

Dung of cow, buffalo, and goat dung and sewage sludge were collected from different parts of Gorakhpur city. These wastes were exposed to sunlight for 5 to 10 days for the removal of various harmful and noxious gases. *E. fetida* is an epigeic species used for this experiment collected from Vermibiotechnology Laboratory, Department of Zoology, D.D.U., Gorakhpur University, Gorakhpur. In the laboratory condition, the temperature was 20 °C to 30 °C and the moisture was maintained up to 40% to 60% for proper growth and survival of earthworms (Bhartiya and Singh, 2011; Nath and Singh, 2016).

### Experimental setup for vermicomposting

Vermicomposting of different combinations of animal (cow, buffalo, and goat) dung with sewage sludge in 1:1, 1:2, and 1:3 ratios was used for the preparation of vermibeds which was conducted on the cemented earth surface. The size of each vermibed is (3 m length × 1 m width × 12 cm depth). 2 kg of cultured *E. fetida* inoculated in each bed and covered with jute pockets for proper moisture maintenance up to 40 to 50 days. Each vermibed manually turned over up to 5 weeks after a one-week interval. After 60 days, tea granules like vermicompost appeared on the upper surface of each bed. The prepared vermicompost and earthworms were used for chemical analysis and heavy metals Pb, Cr, and Cd concentrations (Bhartiya and Singh, 2012b; Nath and Singh, 2016).

### Analysis of heavy metals in initial feed mixture, final vermicompost, and earthworm body

Heavy metal content of the initial feed mixture (IFM) and final vermicompost (FVC) was measured by the method of Maboeta and Rensburg (2003). 1 g of IFM and FVC is required for the preparation of samples. These samples were digested by adding nitric acid (1:1) placing on a hot plate and heating it for 4 hours at 90 °C to 100 °C and keep watching that the sample did not dry up after digestion sample was poured into a 100 mL flask filter through Whatman No 41 paper.

The heavy metals in the earthworm's body tissue are digested using the method of Katz and Jenneis (1983). Earthworms dried, ground, and burned to ash at high temperatures. The ash was placed in a test tube with about 10 mL to 15 mL of 55% nitric acid (HNO<sub>3</sub>) added to it. The solution was left for 12 hrs at room temperature. The solution heated at 40 °C to 60 °C for 2 hours and followed by 120 °C to 130 °C for one hour and 1 mL of 70% perchloric acid (HClO<sub>4</sub>) was added to it. The solution was allowed to cool and 5 mL of distilled water was added thereafter. The solution was again heated up to 130 °C until white fumes emitted. The solution was allowed to cool and filtered through the Whatman No 41 filter paper. This solution allowed for the analysis of heavy metals of the initial feed mixture, final vermicompost, and earthworm body by Shimadzu AA-7000 atomic absorption spectrometer (Najah et al., 2015).

### Calculation of heavy metals concentration in vermicompost and earthworm body

$$\% \text{decrease HMCs in VC} = \frac{(\text{IFM} - \text{FVC})}{\text{IFM}} \times 100 \text{ mg/kg}$$

$$\% \text{increase HMCs in earthworm body} = \frac{(\text{EfBFV} - \text{EfBC})}{\text{EfBC}} \times 100 \text{ mg/kg}$$

HMCs = heavy metals concentration

IFM = initial feed mixture

FVC = final vermicompost

EfBC = earthworm *Eisenia fetida* body (control)

EfBFV = earthworm *Eisenia fetida* body in final vermicompost

### Statistical analysis

All the data are mean ± SD of 6 replicates. Students 't' test was applied to determine the significant (P < 0.05) difference between combinations of cow, buffalo, and goat dung with sewage sludge (SS) of initial, final vermicompost, before inoculation and final earthworm *E. fetida* body (Sokal and Rohlf, 1973).

## 3. Results and discussion


Data emerging from results show there was a significant reduction of Pb, Cr, and Cd observed in prepared vermicompost of different animal dung and sewage sludge in the ratios 1:1, 1:2, and 1:3. Chromium (Cr) was significantly decreased in cow dung (CD) and goat dung (GD) and cadmium (Cd) in CD, CD+SS (1:1), BD and GD at BDL level.

The Lead (Pb) was significantly decreased by 76.470% ( $0.527 \pm 0.004$  mg/kg to  $0.124 \pm 0.003$  mg/kg) in the goat dung with sewage sludge GD+SS 1:1 ratio (Table 1 and Fig. 1), whereas Pb accumulated and significantly increased ( $5.248 \pm 0.007$  mg/kg to  $6.456 \pm 0.009$  mg/kg) 23.018% in earthworm *Eisenia fetida* body in the combination of buffalo dung with sewage sludge (BD+SS, 1:3). The Cr and Cd were significantly increased in the *E. fetida* body 3.384% ( $11.435 \pm 0.012$  mg/kg to  $11.822 \pm 0.009$  mg/kg) and 11.801% ( $3.347 \pm 0.008$  mg/kg to  $3.742 \pm 0.010$  mg/kg), respectively, in the combination of buffalo dung with sewage sludge (BD+ SS) in ratio 1:3 (Table 2 and Fig. 2). Khwairakpam and Bhargava (2009) reported that the combination of metal decreased during vermicomposting with all worm with *Eudrilus euginae*, *Perionyx excavates* and *Eisenia fetida* slightly more in case of lead.


*Eisenia fetida* had a significant degree of accumulation of Cr, Pb, and Zn compared to *Eudrilus euginae*, and the bioaccumulation factor was higher in the case of *E. fetida* (Parseh et al., 2021). The bioaccumulation of metals in the body of *E. fetida* and *E. andrei* was thus  $Cd > Zn > Cu > Ni > Pb$  (Rorat et al., 2016). The metallothioneins (MTs)—

small, cysteine- rich proteins of *Lumbricus rubellus* and common red earthworm (*Eisenia fetida*) that bind d10 metal ions (Zn(II), Cd(II), or Cu(I)) in clusters. Three MT homologues are known for one of which, wMT- 2, is strongly induced by exposure of worms to cadmium (Kowald et al., 2016). Earthworms have the ability to accumulate heavy metals, including Cu, Cd, Pb, and Zn by earthworm occurs through two pathways, which include absorption following dermal contact and adsorption through the intestinal tissues (Nirola et al., 2016; Wang et al., 2018; Lanno et al., 2019; Srut et al., 2019).

There was a significant increase of heavy metals in the body of inoculated earthworm *Eisenia fetida* after vermicomposting in a combination of animal dung with biological wastes during vermic- activity (Shahmansouri et al., 2005; Jordao et al., 2006; Bhartiya and Singh, 2012a, 2012b; Singh and Bhartiya, 2020a). Significant changes of fraction distribution and bioavailability of these metals (Zn, Pb, Fe, Mn, Cr, Co, and Cu) occurred during earthworm activity (De- vlieghe and Verstraete, 1996). A significant concentration of Cd accumulation was reported in the earthworm body of *Eisenia fetida* (Conder et al., 2002; Shahmansouri et al., 2005; Singh et al., 2014; Singh and Bhartiya, 2020b).



**Figure 1.** Concentration of heavy metals (mg/kg) in combination of different animal dung with sewage sludge in initial feed mixture and vermicompost.



**Figure 2.** Concentration of heavy metals (mg/kg) of earthworm *Eisenia fetida* body in different of combination animal dung with sewage sludge after vermicomposting.

**Table 1.** Concentration of heavy metals (mg/kg) in a combination of different animal dung with sewage sludge and vermicompost.

Particulars	Ratio	Heavy metal concentration in mg/kg								
		Pb			Cr			Cd		
		IFM	FVC	% Decrease	IFM	FVC	% Decrease	IFM	FVC	% Decrease
SS	-	2.196±0.010	-	-	0.794±0.006	-	-	0.538±0.009	-	-
CD	-	0.331±0.005	0.105±0.003*	68.278	0.199±0.004	BDL*	100	0.045±0.005	BDL*	100
CD+SS	1:1	0.530±0.003	0.297±0.005*	43.963	0.356±0.005	0.085±0.003*	76.123	0.209±0.005	BDL*	100
	1:2	0.905±0.005	0.340±0.004*	62.431	0.467±0.004	0.162±0.004*	65.310	0.381±0.004	0.112±0.006*	70.603
	1:3	1.560±0.008	0.890±0.004*	42.948	0.598±0.004	0.178±0.006*	70.234	0.390±0.005	0.129±0.005*	66.923
BD	-	0.452±0.004	0.119±0.005*	73.672	0.236±0.003	0.092±0.004*	61.016	0.062±0.004	BDL*	100
BD+SS	1:1	0.702±0.006	0.255±0.004*	63.675	0.367±0.005	0.156±0.005*	57.493	0.173±0.003	0.043±0.004*	75.144
	1:2	1.025±0.006	0.368±0.006*	64.097	0.402±0.003	0.189±0.003*	52.985	0.210±0.004	0.085±0.005*	59.523
	1:3	1.614±0.008	0.795±0.005*	50.743	0.678±0.003	0.308±0.004*	54.572	0.295±0.007	0.106±0.004*	64.067
GD	-	0.149±0.004	0.055±0.002*	63.087	0.127±0.004	BDL*	100	0.196±0.004	BDL*	100
GD+SS	1:1	0.527±0.004	0.124±0.003*	76.470	0.250±0.005	0.084±0.006*	66.400	0.309±0.003	0.068±0.003*	77.993
	1:2	0.851±0.004	0.209±0.005*	64.027	0.603±0.003	0.196±0.005*	67.495	0.371±0.006	0.096±0.004*	74.123
	1:3	1.272±0.009	0.423±0.006*	66.745	0.675±0.003	0.205±0.008*	69.629	0.467±0.006	0.145±0.005*	68.950

\* Significant P < 0.05 “t” test between animal dung, sewage sludge, and final vermicompost - Each value is the Mean ±SD of six replicates BDL-Below detectible limit i.e. 0.0001 mg/kg  
 SS-sewage sludge, IFM- initial feed mixture, FVC- final vermicompost, CD-cow dung, BD-buffalo dung, GD-goat dung

**Table 2.** Concentration of heavy metals (mg/kg) of earthworm *Eisenia fetida* body in different combinations of animal dung with sewage sludge after vermicomposting.

Particulars	Ratios	Heavy metal concentration in earthworm <i>Eisenia fetida</i> body (mg/kg)								
		Pb			Cr			Cd		
		IFM	FVC	% Decrease	IFM	FVC	% Decrease	IFM	FVC	% Decrease
CD	-	5.248±0.007	5.445±0.006*	3.753	11.435±0.012	11.552±0.008*	1.023	3.347±0.008	3.552±0.005*	6.124
CD+SS	1:1	5.248±0.007	5.406±0.004*	3.010	11.435±0.012	11.659±0.006*	1.958	3.347±0.008	3.565±0.005*	6.513
	1:2	5.248±0.007	5.619±0.003*	7.069	11.435±0.012	11.806±0.006*	3.244	3.347±0.008	3.704±0.004*	10.666
	1:3	5.248±0.007	6.040±0.008*	15.091	11.435±0.012	11.822±0.009*	3.384	3.347±0.008	3.742±0.010*	11.801
BD	-	5.248±0.007	5.662±0.004*	7.888	11.435±0.012	11.588±0.004*	1.337	3.347±0.008	3.411±0.007*	.912
BD+SS	1:1	5.248±0.007	5.769±0.005*	9.927	11.435±0.012	11.592±0.005*	1.372	3.347±0.008	3.463±0.006*	3.465
	1:2	5.248±0.007	6.045±0.003*	15.186	11.435±0.012	11.685±0.006*	2.186	3.347±0.008	3.475±0.005*	3.824
	1:3	5.248±0.007	6.456±0.009*	23.018	11.435±0.012	11.762±0.006*	2.859	3.347±0.008	3.650±0.008*	9.052
GD	-	5.248±0.007	5.450±0.003*	3.849	11.435±0.012	11.505±0.007*	0.612	3.347±0.008	3.501±0.004*	4.601
GD+SS	1:1	5.248±0.007	5.601±0.004*	6.726	11.435±0.012	11.560±0.005*	1.093	3.347±0.008	3.545±0.006*	5.915
	1:2	5.248±0.007	5.670±0.005*	8.041	11.435±0.012	11.723±0.008*	2.518	3.347±0.008	3.673±0.005*	9.740
	1:3	5.248±0.007	6.186±0.005*	17.873	11.435±0.012	11.775±0.009*	2.973	3.347±0.008	3.705±0.007*	10.696

\* Significant  $P < 0.05$  "t" test between earthworm body of each vermibeds before inoculation and after vermicomposting Each value is the Mean  $\pm$ SD of six replicates

SS-sewage sludge, EfBC- earthworm *Eisenia fetida* body (control)

EfBFV- earthworm *Eisenia fetida* body in final vermicomposting, CD-cow dung, BD-buffalo dung, GD-goat dung

## 4. Conclusion

*Eisenia fetida* accumulated heavy metals in their body tissues from different combinations of cow, buffalo, and goat dung with sewage sludge in the ratios 1:1, 1:2, and 1:3. There was a significant decrease in the concentration of Pb, Cr, and Cd in different animal dung with sewage sludge during vermicomposting. Cr and Cd were significantly decreased in cow dung (CD) and goat dung (GD) at the BDL level whereas the Pb concentration significantly decreased (89.261%) in the combination of GD+SS (1:1). *E. fetida* was a suitable species for the accumulation of Pb, Cr, and Cd from different waste during vermicomposting. The result of the study showed that earthworms play an important role in the remediation of heavy metals from sewage sludge with different combinations of animal dung.

## Acknowledgment

The authors are thankful to Dr Jaswinder Singh, Associate Professor, Post Graduate Department of Zoology, Khalsa College, Amritsar, Punjab, India for technical support.

### Author contribution

All authors are contributed in experiment as well as paper drafting and writing.

### Availability of data and materials

The data that support the findings of this study are available from the corresponding author upon reasonable request.

### Conflict of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Open access

This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the OICC Press publisher. To view a copy of this license, visit <https://creativecommons.org/licenses/by/4.0>.

## References

- Ahmed M, Qadeer U, Ahmed ZI, Fayyaz-ul H (2016) Improvement of wheat (*Triticum aestivum*) drought tolerance by seed priming with silicon. *Archi Agron Soil Sci* 62 (3): 299–315.
- Bhartiya DK, Singh K (2011) Accumulation of heavy metals by *Eisenia fetida* from different animal dung and kitchen wastes during vermicomposting. *Int J Life Sci Technol* 4 (7): 47–52. <https://doi.org/10.1080/03650340.2015.1048235>
- (2012a) Heavy Metals accumulation from municipal solid wastes with different animal dung through vermicomposting by earthworm *Eisenia fetida*. *World Appl Sci J* 17 (1): 133–139.

- (2012b) Heavy metals remediation from maize (*Zea mays*) crop by the use of vermicomposts through vermicomposting by *Eisenia fetida*. *American-Eurasian J Agric Environ Sci* 12 (9): 1215–1222. <https://doi.org/10.5829/idosi.ajeaes.2012.12.09.1678>
- Cleveland LM, Minter ML, Cobb KA, Scott AA, German VF (2008) Lead hazards for pregnant women and children. *Am J Nurs* 108:40–49. <https://doi.org/10.1097/01.NAJ.0000337736.76730.66>
- Conder JM, Seals LD, Lanno RP (2002) Method for determining toxicologically relevant cadmium residues in the earthworm *Eisenia fetida*. *Chemosphere* 49:1–7. [https://doi.org/10.1016/s0045-6535\(02\)00192-3](https://doi.org/10.1016/s0045-6535(02)00192-3)
- Devliegher W, Verstraete W (1996) *Lumbricus terrestris* in a soil core experiment: Effect of nutrient- enrichment processes (NEP) and gut associated processes (GAP) on the availability of plant nutrients and heavy metals. *Soil Biol Biochem* 28:489–496. [https://doi.org/10.1016/0038-0717\(95\)00190-5](https://doi.org/10.1016/0038-0717(95)00190-5)
- Dume B, Hanc A, Svehla P, Michal P, Chane AD (2022) Vermicomposting technology as a process able to reduce the content of potentially toxic elements in sewage sludge. *Agronomy* 12 (9): 1–15. <https://doi.org/10.3390/agronomy12092049>
- Garg P, Gupta A, Satya S (2006) Vermicomposting of different types of waste using: A comparative study. *Bioresur Technol* 97:391–395. <https://doi.org/10.1016/j.biortech.2005.03.009>
- Gupta PK (2005) Vermicomposting for sustainable agriculture. *Bharat Printing Press, Jodhpur*, 11–14.
- Iwai CB, Ta-Oun M, Chuasavatee T, Boonyotha P (2013) Management of municipal sewage sludge by vermicomposting technology: Converting a waste into a bio fertilizer for agriculture. *Int J Environ Rural Dev* 4:169–174.
- Jarup L (2003) Hazards of heavy metal contamination. *Br Med Bull* 68 (1): 167–182. <https://doi.org/10.1093/bmb/ldg032>
- Jatav HS, Singh SK, Jatav SS, Latore AM, Kumar V, Singh P (2021) Sewage sludge quality assessment of sewage treatment plant, Bhagwanpur, Varanasi and its safe utilization in agriculture. *J Environ Biol* 42:512–517. [https://doi.org/10.22438/jeb/42/2\(SI\)/SI-271](https://doi.org/10.22438/jeb/42/2(SI)/SI-271)
- Jordao CP, Nascentes CC, Cecon PR, Fontes RLF, Pereira JL (2006) Heavy metal availability in Soil amended with composted urban solid wastes. *Environ Monit Asses* 112:309–326. <https://doi.org/10.1007/s10661-006-1072-y>
- Katz SA, Jenneis SW (1983) Regulatory compliance monitoring by atomic absorption spectroscopy. *Verlay Chemical Int, Fl*
- Kaviraj, Sharma S (2003) Municipal solid wastes management through vermicomposting employing exotic and local species of earthworm. *Bioresur Technol* 90:169–173. [https://doi.org/10.1016/S0960-8524\(03\)00123-8](https://doi.org/10.1016/S0960-8524(03)00123-8)
- Khwairakpam M, Bhargava R (2009) Vermitechnology for sewage sludge recycling. *J Hazard Mater* 161:948–954. <https://doi.org/10.1016/j.jhazmat.2008.v04.088>
- Kowald GR, Sturzenbaum SR, Blindauer CA (2016) Earthworm *Lumbricus rubellus* MT-2: metal binding and protein folding of a true Cadmium-MT gregory. *Int J Mol Sci* 17 (1): 65. <https://doi.org/10.3390/ijms17010065>
- Lanno RP, Oorts K, Smolders E, Albanese K, Chowdhury MJ (2019) Effects of soil properties on the toxicity and bioaccumulation of lead in soil invertebrates. *Environ Toxicol Chem* 38:1486–1494.
- Lee WK, Abouhamed M, Thevenod F (2006) Caspase-dependent and -independent pathways for cadmium-induced apoptosis in cultured kidney proximal tubule cells. *Am J Physiol Renal Physiol* 291 (4): 823–832. <https://doi.org/10.1152/ajprenal.00276.2005>
- Li X, Xing M, Yang J, Huang Z (2011) Compositional and functional features of humic acid-like fractions from vermicomposting of sewage sludge and cow dung. *J Hazard Mater* 185:740–748. <https://doi.org/10.1016/j.jhazmat.2010.09.081>
- Maboeta MS, Rensburg LV (2003) Vermicomposting of industrially produced wood chips and sewage sludge utilizing *E. foetida*. *Ecotoxicol Environ Saf* 56:265–270. [https://doi.org/10.1016/S0147-6513\(02\)00101-X](https://doi.org/10.1016/S0147-6513(02)00101-X)
- Najah Z, Elsherif KM, Alshtewi M, Attorshi H (2015) Phytochemical profile and heavy metals contents of *Codium tomentosum* and *Sargassum hornschurchi*. *J Appl Chem* 4 (6): 1821–1827.
- Nath G, Singh K (2016) Vermiwash: A potent liquid biofertilizer. *Res J Sci Technol* 8 (1): 21–30. <https://doi.org/10.5958/2349-2988.2016.00003.6>
- Nath G, Singh K, Singh DK (2009) Chemical analysis of vermicomposts/vermiwash of different combinations of animal, agro and kitchen wastes. *Aust J Bas Appl Sci* 3 (4): 3671–3676.
- Nirola R, Megharaj M, Saint C, Aryal R, Thavamani P, Venkateswarlu K, Naidu R, Beecham S (2016) Metal bioavailability to *Eisenia fetida* through copper mine dwelling animal and plant litter, a new challenge on contaminated environment remediation. *Int Biodeter Biodegrad* 113:208–216. <https://doi.org/10.1016/j.ibiod.2016.03.007>
- Panday R, Basnet BB, Bhatt PS, Tamrakar AS (2014) Bio-concentration of heavy metals in vermicomposting earthworms (*Eisenia fetida*, *Perionyx excavatus* and *Lampito mauritii*) in Nepal. *J Microbiol Biotechnol Food Sci* 3:416–418.

- Parseh I, Mousavi K, Badienejad A, Mehdi M, Mofrad G, Hashemi M, Azadbakht O, Karimi H (2021) Microbial and composition changes during vermicomposting process resulting from decomposable domestic waste, cow manure and dewatered sludge. *Int J Env Health Eng* 10:3. [https://doi.org/10.4103/ijehe.ijehe\\_56\\_20](https://doi.org/10.4103/ijehe.ijehe_56_20)
- Prajapati UB, Singh S (2013) Analysis of sewage in peri-urban region of gorakhpur-U.P., India. *J Water Res Photon* 135:257–261.
- Purdea L, Rusanescu CO, Tucureanu MC (2019) Alternative for the use of sewage sludge in Romania. *Rev Chim* 70:1967–1970. <https://doi.org/10.37358/RC.19.6.7255>
- Rorat A, Suleiman H, Grobelak A, Grosser A, Kacprzak M, Plytycz B, Vandenbulcke F (2016) Interactions between sewage sludge-amended soil and earthworms-comparison between *Eisenia fetida* and *Eisenia andrei* composting species. *Env Sci Pollut Res* 23:3026–3035. <https://doi.org/10.1007/s11356-015-5635-8>
- Rusanescu CO, Rusanescu M, Voicu G, Paraschiv G, Biris SS, Popescu IN (2022) The recovery of vermicompost sewage sludge in Agriculture. *Agronomy* 12 (2653): 1–12. <https://doi.org/10.3390/agronomy12112653>
- Shahmansouri MR, Pourmoghadas H, Parvaresh AR, Alidadi H (2005) Heavy metals bioaccumulation by Indian and Australian earthworm (*Eisenia fetida*) in the sewage sludge vermicomposting. *Iranian J Env Health Sci Eng* 2 (1): 28–33.
- Singh K, Bhartiya DK (2021) A review article on vermibiotechnology and waste management. *World Rev Sci Technol Sust Dev* 17 (1): 81–97. <https://doi.org/10.1504/WRSTSD.2021.113976>
- (2020a) Heavy metal accumulation by earthworm *Eisenia fetida* from animal waste, soil and wheat (*Triticum aestivum*) for protection of human health. *Res J Pharm Technol* 13 (7): 3205–3210. <https://doi.org/10.5958/0974-360X.2020.00567.3>
- (2020b) Minimization of health problems through vermibiotechnology and accumulation of heavy metal from biological waste, soil and rice (*Oryza sativa* L.). *Res J Pharm Technol* 13 (8): 3580–3586. <https://doi.org/10.5958/0974-360X.2020.00633.2>
- Singh K, Bhartiya DK, Chauhan HK, Rai R, Singh RN (2014) bioaccumulation of cobalt and lead by earthworm *eisenia fetida* from sewage sludge with different cattle dung during vermicomposting. *Res J Sci Technol* 6 (4): 175–179.
- Sokal RR, Rohlf FJ (1973) Introduction to biostatistics. *W.H. Freeman Company*, 363.
- Spurgeon DJ, Hopkin SP (1999) Comparisons of metal accumulation and excretion kinetics in earthworms (*Eisenia fetida*) exposed to contaminated field and laboratory soils. *Appl Soil Ecol* 11:227–243. [https://doi.org/10.1016/S0929-1393\(98\)00150-4](https://doi.org/10.1016/S0929-1393(98)00150-4)
- Srut M, Menke S, Hockner M, Sommer S (2019) Earthworms and cadmium- heavy metal resistant gut bacteria as indicators for heavy metal pollution in soils. *Ecotoxicol Environ Saf* 171:843–853. <https://doi.org/10.1016/j.ecoenv.2018.12.102>
- Suleiman MS, Wasonga VO, Mbau JS, Suleiman A, Elhadi YA (2017) Non-timber forest products and their contribution to households' income around Falgore Game Reserve in Kano, Nigeria. *Ecol Process* 6 (23): 1–14. <https://doi.org/10.1186/s13717-017-0090-8>
- Suthar S (2006) Potential utilization of guar gum industrial waste in vermicompost production. *Bioresur Technol* 97:2474–2477. <https://doi.org/10.1016/j.biortech.2005.10.018>
- Wang K, Qiao Y, Zhang H, Yue S, Li H, Ji X, Liu L (2018) Bioaccumulation of heavy metals in earthworms from field contaminated soil in a subtropical area of China. *Ecotoxicol Environ Saf* 148:876–883. <https://doi.org/10.1016/j.ecoenv.2017.11.058>
- WHO (1993) Evaluation of Certain Food Additives and Contaminants: Forty-First Report of the Joint FAO/WHO Expert Committee on Food Additives. *WHO Technical Report Series; Geneva, Switzerland*, 837.
- (1985) GEMS: Global Environmental Monitoring System, Guidelines for the Study of Dietary Intakes of Chemical Contaminants. *WHO Offset Publication; Geneva, Switzerland*, 87.
- Yadav A, Garg VK (2011) Industrial waste and sludge management by vermicomposting. *Rev Environ Sci Biotechnol* 10 (3): 243–276. <https://doi.org/10.1007/s11157-011-9242-y>
- Yadav KD, Tare V, Ahammed MM (2012) Integrated composting–vermicomposting process for stabilization of human faecal slurry. *Ecol Eng* 47:24–29. <https://doi.org/10.1016/j.ecoleng.2012.06.039>