







# A comprehensive review on organic waste compost as an effective phosphorus source for sustainable agriculture

Lady Johanna Bohórquez-Sandoval<sup>1</sup> , José Francisco García-Molano<sup>2,\*</sup> ,  
José Antonio Pascual-Valero<sup>3</sup> , Margarita Ros-Muñoz<sup>3</sup> 

<sup>1</sup>University of Murcia, International Doctoral School, University Campus of Espinardo, Murcia, Spain.

<sup>2</sup>Juan de Castellanos University Foundation, Department of Agricultural and Environmental Sciences Tunja, Colombia.

<sup>3</sup>CSIC-CEBAS. Department of Soil and Water Conservation, University Campus of Espinardo, Murcia, Spain.

\*Corresponding author: [jgarcia@jdc.edu.co](mailto:jgarcia@jdc.edu.co)

## Review Paper

Received:

05 October 2023

Revised:

13 February 2024

Accepted:

20 June 2024

Published online:

07 August 2024

© The Author(s) 2024

## Abstract:

**Purpose:** The transformation of Organic Waste through composting has become a source of organic matter that, through microbial action, generates molecules conducive to phosphate solubilization and produces fractions of Po and Pi readily available for plants. For this reason, this review focused on analyzing various research studies where Organic Waste is utilized to release available phosphorus and thus improve agricultural production.

**Method:** One hundred and eighteen articles were analyzed and synthesized from recognized databases, addressing the following topics: Phosphorus in the Earth's crust, phosphorus recovery from Organic Waste, compost, metabolic pathways for phosphorus release in compost, and microbial activity for phosphorus release in compost.

**Results:** The selection of organic materials and the incorporation of MSP during the composting process favor the availability of P for plants. The compost microbiota associated with phosphorus not only enhances its availability but also may promote plant growth. Various mechanisms exist for P release, such as the enzymes  $\beta$ -1,4-glucosidase and  $\beta$ -D-fructofuranosidase, along with mycorrhizae facilitating P transport, the enzymatic cofactor pyrroloquinoline quinone, and the function of phosphorus genes contributing to its availability, thereby promoting sustainable agricultural practices.

**Conclusion:** Composts can serve as valuable P sources for plants, wherein their existing microbiota enhances phosphorus availability, facilitating increased phosphorus recycling through symbiotic relationships, microbial interactions, and the action of phosphorus genes, thereby ensuring sustainable agriculture.

**Keywords:** Enzymatic activity; Organic acids; Organic waste; Phytohormones; Phosphate solubilizers

## 1. Introduction

Phosphorus plays a vital role in various physiological functions in plants, such as cell division, root development, and photosynthesis, as it is a component of essential molecules like phospholipids, nucleic acids, and ATP (Ortega, 2020). However, despite the assistance of specialized microorganisms in releasing phosphorus from compounds, it remains relatively scarce in the soil for plants (Atoloye et al., 2021). Phosphorus (P) is the eleventh most abundant element in the Earth's crust (Corbridge, 2013), with a gross abundance ranging from 0.10% to 0.12% (w/w) (Fuller, 1972; Cor-

bridge, 2013). Therefore, total mineral phosphorus concentrations in soil range from 35 to 5300 mg P/Kg, with an average of approximately 800 mgP/Kg to 1400 mgP/kg (Bowen, 1979; Azam et al., 2019; Walton et al., 2023). Mineral phosphorus in soil originates from igneous and sedimentary rocks, which are initially insoluble and must be solubilized for plant assimilation (Contrato, 2018). This input has been excessive over the years through highly soluble phosphates, and the unused remnants accumulate in the soil, either through multivalent metal chelation or clay and organic matter absorption, requiring application in each crop cycle. High concentrations of phosphate can

be found bound to other elements such as Ca, F, or Cl, forming apatite and fluorapatite,  $(Ca_5(PO_4)_3F)$ , chlorapatite  $(Ca_5(PO_4)_3Cl)$ , and hydroxyapatite  $(Ca_5(PO_4)_3OH)$  (Espinel, 2020). Other mineral sources of P include lazulite  $(Mg-Fe)$ ,  $Al_2(PO_4)_2(OH)_2$ , strengite  $(FePO_4 \cdot 2H_2O)$ , vauxite  $(FeAl_2(PO_4)_2(OH)_2 \cdot 6H_2O)$ , vivianite  $(Fe_3(PO_4)_2 \cdot 8H_2O)$ , and variscite  $(AlPO_4 \cdot 2H_2O)$  (Veith and Sposito, 1977).

Approximately 30% to 65% of total soil phosphorus exists in organic forms, with the remainder in inorganic forms. Organic forms consist of dead plant/animal residues and soil microorganisms, which are crucial for converting them into plant-usable forms (Tapia-Torres and García-Oliva, 2013). However, soil microorganisms may hinder phosphorus absorption through mineralization processes (Kwesi-Asomaning, 2020; Singh et al., 2015).

Phosphate rocks, essential for phosphorus compounds and fertilizers, are globally transported at a rate of approximately 30 million tons annually. Morocco, with 85% of its reserves, extracts phosphorus from alkaline intrusive igneous rocks like apatite (Contrato, 2018; Azam et al., 2019). Despite finite resources, with about 100 years of reserves left, concerns arise about potential hoarding by major producers and its impact on global food security (Escamilla, 2015). Apatite extraction may introduce high levels of heavy metals and radioactive elements, posing risks to biomass and disrupting carbon and nitrogen cycles (Basílio et al., 2022).

Recycling organic materials, such as sewage sludge, Organic Waste, invasive weeds, and agro-industrial residues, offers an alternative phosphorus source (Grigatti et al., 2017; Oliveira et al., 2019; Kauser and Khwairakpam, 2022; Hernández-Lara et al., 2021; Zuhair et al., 2022). Studies show that composted organic materials provide superior soil-available phosphorus compared to chemical fertilizers (Adnan et al., 2017). Targeted recycling of Organic Waste and sludge is crucial for reintroducing nutrients into the soil, enhancing biofertilizer production, increasing agricultural output, and addressing public health, environmental protection, and climate change concerns (Galvis, 2016; Hernández-Berriel et al., 2016; Vergara and Tchobanoglous, 2012).

This review stems from the research question: What molecules present in organic matter wastes have the potential to solubilize phosphates and how can they be managed through composting and its relationship to agricultural production sustainability? For this reason, the primary objective of this work is to identify molecules present in organic matter that possess properties for phosphate solubilization and to explore the composting process of Organic Waste aiming to leverage the phosphorus (P) release capacity as an essential element for plant growth.

In a world where phosphorus availability is crucial for agricultural production, understanding how these molecules can influence phosphate solubilization could be key to improving the efficiency and sustainability of fertilization systems. In this way, it could not only contribute to optimizing the use of natural resources but also to reducing the reliance on synthetic fertilizers and their potential environmental

impacts.

## 2. Materials and method

One hundred and eighteen articles were obtained from our search, which was conducted following the methodology proposed by Ferenhof and Fernandes (2016). We used Scopus, Science Direct, Web of Science, PubMed, Scielo, Google Scholar, ResearchGate, and Dialnet databases and considered articles published from 1972 to 2023. The search focused on the following keywords or topics: Phosphorus in the Earth's crust, phosphorus recovery from Organic Waste, compost, metabolic pathways for phosphorus release in compost, and microbial activity for phosphorus release in compost.

### The need for new sources of Phosphorus in agriculture

Applying P to crops in their assimilable forms leads to better crop yields (Moreno and Moral, 2007). However, excessive application through synthetic fertilizers can have adverse effects on its availability since elements such as aluminum (Al) and iron (Fe) in acidic soils or calcium (Ca) in basic soils can sequester P, resulting in significant P loss (Chen et al., 2022). Excessive application also increases P loss through leaching and migration from agricultural soils, leading to eutrophication and subsequent anoxia in marine ecosystems. Therefore, P is prevented from re-circulating, affecting its potential to absorb  $CO_2$  (Fernández-Marcos, 2011; Iida and Shock, 2011; Richardson et al., 2023). Furthermore, higher prices for phosphate fertilizers due to rising demand and input costs raise production expenses for farmers (Moharana et al., 2020). The above-mentioned reasons have encouraged us to find new P sources and the optimization to its application to optimize its plant uptake.

### Phosphorus recovery from organic waste and sewage sludge

An increase in human activity has resulted in a greater variety of waste compositions, with approximately 46% of solid waste being organic, primarily from food preparation (Hoang et al., 2022), where improper management of this Organic Waste can lead to potential contamination (Bank, 2018). Organic Waste (OW) and urban and agro-industrial sewage sludge not only contain abundant organic matter (>40%) but also serve as significant nutrient sources. Sustainable management can valorize these waste materials as nutrients for soil and plants. The OW category includes the organic fraction of municipal solid waste, wood chip waste, agricultural waste, garden waste, food waste, and animal manure, with variations depending on their origin (Table 1). The phosphorus content of the reviewed OW ranges from 0.05% to 1.78%, indicating relatively low phosphorus values (Moharana et al., 2020; Wei et al., 2018b). In contrast, sewage sludge exhibits higher total phosphorus values (1.53% to 1.78%) (Grigatti et al., 2017) (Table 1). This higher content is likely attributed to soap residues containing phosphorus (Djandja et al., 2022) and residues from food such as legumes, vegetables, cereals, tubers, fruits, and bone waste in the form of organophosphonates (Wei et al., 2018b; Tapia-Torres and García-Oliva, 2013). Apart

from phosphorus, both OW and sewage sludge contain other essential nutrients for plant growth, including nitrogen, ranging from 0.21% to 3.90%. These nutrients originate from protein-rich materials, amino acids, some B-complex vitamins, and Phytohormones found in fruits, vegetables, and forage. These materials act as food for decomposer microorganisms, thereby stimulating the proliferation of free-living nitrogen-fixing microorganisms. Some of these microorganisms, such as *Aeromonas salmonicida*, *Pausterella pneumotropica*, *Burkholderia tropica*, and *Bacillus* sp., possess dual functions as they not only fix nitrogen but also solubilize phosphate (Bolívar-Anillo et al., 2016; Corrales et al., 2014; Pérez-Cordero et al., 2014). However, lower nitrogen values can increase the C/N ratio, limiting microbial activity in Organic Waste management (Santamaria and Ferrera, 2002).

Another essential nutrient for plant growth is potassium, which plays a role in root cell growth, improving plant nutrient uptake. Potassium content ranges from 0.95% to 2.32% (Moharana et al., 2020; Pedrosa et al., 2013; Pascual et al., 1999), although data on potassium do not usually appear in the reviewed studies. Kitchen waste shows high levels of K, probably caused by banana peels, which are known for their high potassium content. This Organic Waste (OW) and sewage sludge show pH values ranging from 4.83 to 10.2, and they are predominantly basic. Regarding electrical conductivity (EC), they show values below 2, indicating little presence of salts in the raw material, except for MSW (4.90), possibly due to carbonate, sulfate, chloride, or nitrate content (Babana et al., 2013).

### Phosphorus recycling from OW and sewage sludge: Composting

Composting is a microorganism-driven aerobic process that transforms organic materials into stable organic fertilizers (compost). Composts can enhance soil properties, promoting plant growth, mitigate emissions (Hafez et al., 2021; Moreno and Moral, 2007; Zhang et al., 2018) and suppress crop pathogens, (Hernández-Lara et al., 2021). Also, microorganisms, including bacteria and fungi, facilitate phosphorus solubilization during composting (Beltrán-Pineda, 2015), through the production of Organic Acids or enzymes that increase plant-absorbable phosphorus forms (Oliveira et al., 2019). Specifically, microorganisms transform phosphorus by mineralizing organic phosphorus (Po) via phosphatases and producing low molecular weight Organic Acids, primarily by bacteria solubilizing phosphates from inorganic phosphorus (Pi) (Antoun, 2012).

Increasing P availability in compost involves biotechnological manipulation of phosphorus-rich materials, such as manures, phosphate rock, sewage sludge, or biochar during composting (Grigatti et al., 2017; Nobile et al., 2022; Zhang et al., 2018). The use of these materials offers advantages, such as valorizing Organic Waste, reducing production costs (Gao et al., 2019), and enhancing phosphorus solubility in the soil solution, reducing P fixation (Adnan et al., 2017). Also, inoculation during composting with phosphate-solubilizing microorganisms (PSM) boosts phos-

phates in compost, and it is crucial for low molecular weight Organic Acid production and Enzymatic activity linked to phosphorus solubilization-mineralization (Table 2) (Gand, 2014; Moharana et al., 2020; Yadav et al., 2017; Wei et al., 2018b; Zhan et al., 2021).

In addition to composting, vermicomposting a process carried out by earthworms, principally the red Californian earthworm, (*Eisenia fetida* or *Eisenia andrei*) can also transform organic residues into compost. This process is more efficient for biologically treating pre-processed materials like livestock manure (Ferraz et al., 2022) since it promotes the metabolism of easily assimilable molecules, such as simple carbohydrates, peptides, and proteins. It also increases P, Ca, and Mg levels through the mineralization process of the raw materials used.

Composts have regulations to ensure their quality and safety. In the European Union, Regulation (EU) 2019/1009 on solid organic fertilizers establishes the minimum parameter for P at 1 – 2% per mass of P<sub>2</sub>O<sub>5</sub> (EU, 2019). In the specific case of Colombia, solid and liquid organic fertilizers must be endorsed by NTC 5167 (ICONTEC, 2011), which also requires reporting if the phosphorus concentration exceeds 1%, as this affects application to soil or crops.

### The role of compost microbiota in P availability

The availability of phosphorus in both soil and organic residues is guided by phosphorus-solubilizing microorganisms (bacteria (BSP) and fungi (FSP) and *Arbuscular Mycorrhizal Fungi* (AMF)), which are specifically responsible for solubilizing or mineralizing it (Fig. 1, Table 2). The principal BSPs for this function is *Pseudomonas*, *Bacillus*, *Burkholderia*, *Klebsella*, *Agrobacterium*, *Aeromonas*, *Rhizobium*, *Alcaligenes*, *Achromobacter*, and *Lactobacillus* (Table 2), while the most representative groups of FSP are *Aspergillus*, *Penicillium*, *Trichoderma*, and *Paecilomyces* (Table 2). Most of these microorganisms belong to the rhizosphere of plants, but they can also be found in organic matter, which is the main component of organic residues. P solubilization and mineralization are performed by microorganisms through their metabolism, involving the extracellular production of short-chain Organic Acids for inorganic phosphorus or enzymes for organic phosphorus (Fig. 1). Some microorganisms also perform other functions, such as promoting plant growth (Silva et al., 2023). Thus, their presence is valuable for agriculture.

Microorganisms not producing halozone for phosphorus solubilization may employ alternative mechanisms, such as siderophores, which chelate iron atoms from phosphate compounds. These siderophores are present in gram-negative bacteria, fungi, yeasts, and some plants (Aguado et al., 2012). In contrast, Gram-positive bacteria, known for their resilience to adverse conditions, may be more efficient in phosphate solubilization, particularly in withstanding pH changes during composting and in the soil (Corrales et al., 2014).


Arbuscular mycorrhizal fungi (AMF), including *Acaulospora scrobiculata*, *Glomus deserticola*, *Glomus intraradices*, *Glomus versiforme*, *Acaulospora morrowiae*, *Acaulospora spinosa*, *Gigaspora rosea*, and *Rhizophagus*

*intraradices* (Table 2), establish associations in conditions with cations like Al, Fe, or Ca. These associations, found in approximately 80% of plant species, involve AMF extend-

**Table 1.** Chemical characterization of organic waste and sewage sludge.

Origin	Residues	pH	EC (dS/m)	OM %	TOC %	C/N	N %	P %	K %	References
India	OFMSW	7.27	2.06	85.43	40.29	15.20	2.65			(Awasthi et al., 2015)
	WSW	6.54	0.85	94.15	45.18	86.88	0.52	ND	ND	
	AW	7.21	0.86	95.24	44.36	58.36	0.76			
	YW	7.39	0.74	93.14	40.72	45.75	0.89			
CHD				58.92	34.18	25.30	1.35			
MOH	MSW	ND	ND	58.27	33.80	22.10	1.53	ND	ND	(Rana et al., 2018)
PKL (India)				54.99	31.90	27.10	1.10			
Mauritius (Africa)	R1	7.40	0.29	81.37	47.20	27.00	1.70			(Soobhany, 2018)
	R2	7.30	0.26	65.68	38.10	25.40	1.50	ND	ND	
	R3	7.20	0.19	73.09	42.40	27.20	1.60			
Pombal (Brazil)	CB			43.90	25.46	48.96	0.52	0.15	2.32	(Pedrosa et al., 2013)
	PJ	ND	ND	50.17	29.10	32.69	0.89	0.17	0.20	
	PM			43.96	25.49	20.72	1.23	0.18	0.30	
	SM			39.82	23.09	21.57	1.07	0.52	1.50	
Paipa (Colombia)	MOSW	ND	0.002	61.54	35.70	21.40	1.67	ND	ND	(García-Molano et al., 2021)
Ventaquemada (Colombia)	RC	8.18	ND	77.58	45.00	19.90	2.26	ND	ND	(Bohórquez-Sandoval et al., 2020)
United Kingdom	MSW	6.10	4.90	38.70	ND	1.37	1.64	0.45	0.95	(Pascual et al., 1999)
	AL	10.20	2.00	117.40	68.10	31.80	2.14	1.53		
Italy	SS	7.90	1.00	47.70	27.70	6.90	3.90	1.78	ND	(Grigatti et al., 2017)
	GW	6.80	0.93	43.40	25.20	22.50	1.12	0.26		
	KW1	4.99		98.99	57.42	17.08	3.36	0.86		
China	S	7.02	ND	74.63	43.29	47.25	0.91	0.09	ND	(Wei et al., 2017)
	MM	5.40		93.35	54.15	24.59	2.37	0.78		
India	RS			85.85	49.80	94.20	0.53	0.05	1.07	(Moharana et al., 2020)
	WS			90.68	52.60	90.40	0.58	0.06	1.19	
	MS	ND	ND	87.40	50.70	89.30	0.57	0.08	1.33	
	CS			84.13	48.80	64.70	0.75	0.06	1.19	
	TL			95.85	55.60	121.60	0.46	0.05	1.50	
	CD			64.30	37.30	60.20	0.62	0.14	1.11	
Pakistan	SPW	ND	ND	69.80	40.50	28.30	1.46	0.13	ND	(Billah and Bano, 2014)
	KW2	4.83	1.66	72.85	42.26	13.41	3.15	0.46		
China									ND	(Zhan et al., 2021)
	S	5.61	0.09	81.01	46.99	223.70	0.21	0.21		

**EC:** electrical conductivity; **OM:** Organic matter; **TOC:** total organic carbon; **N:** total nitrogen; **P:** total phosphorus; **K:** total potassium **CHD:** Chandigarh, **MOH:** Mohali, **PKL:** Pankhula, **OFMSW:** Organic fraction of municipal solid waste, **WSW:** Wood chip waste, **AW:** Agricultural waste, **YW:** Yard waste, **MSW:** Municipal solid waste, **R1:** Food waste, **R2:** Paper waste, **R3:** Garden waste, **CB:** Banana peelings, **PJ:** Jurema pruning, **PM:** Marmeleiro pruning, **SM:** Sheep manure, **MOSW:** Municipal organic solid waste, **RC:** Rumens content, **AL:** Agro-industrial sewage sludge, **SS:** Urban sewage sludge, **GW:** Green waste, **KW:** Kitchen waste, **S:** Sawdust, **MM:** Mixed materials, **RS:** Rice straw, **WS:** Wheat straw, **MS:** Mustard stubble, **CS:** Chickpea stubble, **TL:** Tree leaves, **CD:** Livestock manure, **SPW:** Simple poultry waste.



**Figure 1.** Mechanisms for obtaining available plant P  
**PSB:** Phosphorus solubilizing bacteria, **PSF:** Phosphorus solubilizing fungi, **AMF:** Arbuscular mycorrhizal fungi.

ing their hyphae into the plant's rhizosphere and penetrating the root cortex, serving various purposes. This includes increased nutrient absorption by the plant, exploration of a larger soil area, enhanced plant biomass, and the provision of organic carbon to AMF. Additionally, AMF contributes to soil stability by producing glomalin and Phytohormones that facilitate P availability to the plant (Maaloum et al., 2020; Etesami et al., 2021; García-Molano et al., 2022; Perea et al., 2019; Velquez et al., 2017; Wahid et al., 2020). This mycorrhizal association depends on phosphate-solubilizing fungi (PSF) and, to a lesser extent, phosphate-solubilizing bacteria (PSB) (Fig. 1), which play a crucial role in facilitating interactions between AMF and plant roots. These interactions involve chemical pathways for P solubilization, accomplished through PSF production of low molecular weight Organic Acids or enzymes and hormone secretion. Furthermore, AMF exude fructose, activating phosphatase genes in PSB and promoting organic phosphorus mineralization (Maaloum et al., 2020; Etesami et al., 2021; Ordoñez et al., 2016). Compost, with organic molecules resembling plant exudates, supports AMF nutrition and growth, facilitating effective P transport to plants. For this reason, when applied to the soil, compost can likely promote mycorrhizal associations, considering its nutritional contributions to AMF (Yang et al., 2018).

### Enzymatic activity (phosphorus mineralization)

Mineralizing organic phosphorus to release orthophosphates occurs when various microbial groups produce extracellular enzymes that synthesize three hydrolytic enzymes: i. phosphonoacetaldehyde hydrolase (phosphonatease), ii. phosphonoacetate hydrolase, iii. phosphonopyruvate hydrolase; iv. C-P lyase breaks the C-P bond, releasing phosphate ion ( $\text{HPO}_4^{2-}$ ) from the organic P source (Kaur, 2020; Tapia-Torres and García-Oliva, 2013). Pyrophosphatase also hydrolyzes pyrophosphates, which are a group of polyphosphates (Darch et al., 2016). Similarly, non-specific phosphatases dephosphorylate P in phosphorylated proteins, and phytases release P from phytic acid (Kaur, 2020). The latter is mainly produced by species of the *Bacillus* and *Enterobacter* genus, which have the advantage of operating within pH ranges of between 3.5 and 7.5 but are more efficient under alkaline pH. Likewise, within the group of

phytases, the  $\beta$ -helix phytase should be highlighted for its exceptional phosphate solubilizing activity (Corrales et al., 2014), along with species from the genera *Pseudomonas* sp., *Burkholderia* sp., *Alcaligenes* sp., and *Aspergillus* sp., which mineralize P by producing phosphatases and phytases (Babana et al., 2013; Behera et al., 2017; Bolívar-Anillo et al., 2016; Cheng and Wan, 2022).

### Organic acids (phosphate solubilization)

Fungi and bacteria play a crucial role in phosphate solubilization, a process influenced by pH, organic matter interactions, and soil physicochemical characteristics (Babana et al., 2013; Darch et al., 2016). The type of organic acid produced depends on pH, with compost pH being alkaline. Adjusting compost pH to 5.0 enhances organic acid and enzyme efficiency (Darch et al., 2016), however, Marra et al. (2015) propose that organic acid production is pH-independent, and it is mediated by mechanisms like proton exclusion, siderophores, and exopolysaccharide production. Phosphate-solubilizing microorganisms (PSM), produce organic acids from macromolecules like carbohydrates, lipids, and peptides through a fermentative pathway like glycolysis (Beltrán-Pineda, 2015; Paredes-Mendoza and Espinosa-Victoria, 2009). *Gluconic acid* is produced by *Penicillium rugulosum* from glucose or sucrose, and *Aspergillus niger* produces citric acid through sucrose fermentation (Pérez-Navarro et al., 2016). *Gluconic acid* releases phosphorus from Ca-P bonding, while *oxalic acid* releases P from Fe-P and Al-P, acting as chelating agents for these cations (Beltrán-Pineda, 2015). Microorganisms like *Pseudomonas* sp. and *Penicillium* sp. produce citric acid, the most efficient organic acid in phosphate solubilization (Babana et al., 2013). Bacteria of the *Bacillus* genus use different pathways, including tricarboxylic acids, to synthesize malic acid or direct glucose oxidation to produce 2-ketogluconic acid (Table 2) (Corrales et al., 2014). Also, certain bacteria, which produce the enzymatic cofactor pyrroloquinoline quinone (PQQ), are known to contribute to the synthesis of organic acids such as gluconic acid, (Sarr et al., 2020; Vera-Cardoso et al., 2017). Fungi present in compost, despite their smaller population, exhibit higher efficiency in P solubilization, attributed to their production of a larger quantity of organic acids compared to bacteria (Babana et al., 2013). On the other hand, the direct action of phosphorus genes on the solubilization of minerals containing it is well established, involving genes such as *gcd*, *ppx*, and *ppa* for  $\text{P}_i$ , and genes like *phoA*, *phnW*, *phoD*, *phnP*, *phnI*, *phnG*, and *phnJ* for  $\text{P}_o$  (Xu et al., 2023).

### Phytohormones

Phosphate solubilization can also be attributed to phytohormone secretion by various microorganisms from the genera *Pseudomonas*, *Bacillus*, *Azotobacter* and *Azospirillum*. These groups are of interest for phosphate solubilization and also have characteristics such as fixing nitrogen, producing phytohormone, and promoting plant growth (Licea-Herrera et al., 2020). The main Phytohormones studied for phosphorus (P) solubilization include auxins, which promote lateral root production for nutrient absorption while inhibiting primary root growth; ethylene, involved in forming ad-

ventitious roots under P deficiency and linked to acid phosphatase regulation (Basílio et al., 2022); and strigolactones, inducing morphological, physiological, and biochemical changes in plants. Strigolactones contribute to increased root biomass, aiding in soil P scavenging through symbiosis with fungi in mycorrhizal formations. Strigolactones may also acidify the environment by releasing protons from root exudates, working in conjunction with phytase and acid phosphatase. However, excess P in plants can negatively impact strigolactone synthesis, reducing dissolved organic carbon exudation (Santoro et al., 2021). Although research on the role of Phytohormones in P solubilization in compost is in its early stages, it highlights another mechanism benefiting organic residues and sludge, not only as a nutrient source for the soil but also as a source of microorganisms, fostering synergistic relationships between microorganisms

and plants.

### Phosphorus fractions in composts from organic waste and sludge

#### Phosphorus forms in organic wastes and sludge composts

Phosphorus in compost can be present in different forms: i. labile, corresponding to orthophosphates that are easily assimilated by plants; ii. moderately available phosphorus, which is bound to metals such as Fe, Al, and Ca and can be solubilized through organic and inorganic acids, along with organic phosphorus as phosphate esters that can be mineralized by enzymes to convert it into labile phosphorus; and iii. occluded phosphorus, which is present in molecules that are challenging to solubilize and mineralize (Takahashi and Dahlgren, 2016; Tinoco-Varela and Bayuelo-Jiménez,

**Table 2.** Phosphorus-solubilizing microorganisms involved in phosphate solubilization, mineralization, and transport

Type	Microorganism	P solubilization mechanisms	References
Bacteria	<i>Aeromonas salmonicida</i>	Indoleacetic acid (IAA)	(Pérez-Cordero et al., 2014)
	<i>Pasteurella pneumotropica</i>	Alkaline phosphatase enzyme	(Behera et al., 2017)
	<i>Alcaligenes sp.</i>		
Bacteria	<i>Burkholderia tropica</i>	Organic acids and phosphatase enzymes.	(Bolívar-Anillo et al., 2016)
		Plant growth promoter (PGP) and siderophore.	(Corrales et al., 2014)
Bacteria	<i>Gluconacetobacter diazotrophicus</i>	Phytohormones IAA. PGP	(Restrepo-Franco et al., 2015)
	<i>Herbaspirillum seropedicae</i>		(Basílio et al., 2022)
	<i>Paenibacillus lautus</i>		(Pérez-Pazos and Sánchez-López, 2017)
	<i>Herbaspirillum sp.</i>		
	<i>Azospirillum sp.</i>		
Bacteria	<i>Azotobacter sp.</i>	PGP Organic acids, phosphatase enzymes, and, in some cases, phytases.	
	<i>Pseudomonas sp.</i>		
	<i>P. aeruginosa</i>		
	<i>Rhizobium,</i>		
	<i>Burkholderia sp,</i>		
	<i>Achromobacter sp</i>		
	<i>Agrobacterium,</i>		
	<i>Aereobacter</i>		
	<i>Flavobacterium</i>		
	<i>Yarrowia,</i>		
	<i>Streptosporangium</i>		
	<i>Erwinia herbicola</i>		
	<i>Proteus</i>		
	<i>Pantoea</i>		
<i>Mycobacterium</i>			
<i>Anthrobacter</i>			
<i>Enterobacter</i>			
Bacteria	<i>Bacillus sp.; B</i>	Organic acids, phosphatase, and phytase enzymes. PGP.	(Estrada-Bonilla et al., 2017)
	<i>liqueniformis</i>		(Yadav et al., 2017)
	<i>B. amyloliquefaciens; B.</i>		(Zhang et al., 2021)
	<i>megaterium; B. firmus</i>		(Zhan et al., 2021)
	<i>Brevibacillus sp</i>		(Adnan et al., 2017)
Bacteria	<i>Lactobacillus sp</i>	Organic acids.	(Estrada-Bonilla et al., 2017)
	<i>Herbiconiux sp</i>		(Zhan et al., 2021)
	<i>Halopolyspora sp</i>		
	<i>Cosenzaea sp</i>		
	<i>Aeromicrobium sp</i>		
	<i>Cellulosimicrobium sp</i>		
Bacteria	<i>Lentibacillus sp</i>		
	<i>Klebsella sp.; K. oxytoca</i>	Mobilization of organic and inorganic P. Organic acids.	(Zhan et al., 2021)
	<i>Micrococcus sp</i>		(Babana et al., 2013)
	<i>Agrobacterium</i>		
<i>tumefaciens</i>			

Bacteria	<i>Firmicutes</i> <i>Proteobacteria</i> <i>Bacteroidetes</i> <i>Bacillus</i> sp.	Organic acids.	(Wei et al., 2017)
Bacteria	<i>B. megaterium</i> <i>B. subtilis</i> <i>B. cereus</i>	Organic acid.	(Castillo-Arteaga et al., 2016) (Panhwar et al., 2013) (Saeid et al., 2018)
Bacteria	<i>Pseudomonas</i> sp, <i>Pantoea</i> , <i>Mycobacterium</i> <i>Bacillus</i> , <i>Rhizobia</i> , <i>Burkholderia</i> , <i>Arthrobacter</i> <i>Enterobacter</i>	Organic acids.	(Adnan et al., 2017)
Bacteria	<i>Rhizobium tropici</i> <i>Acinetobacter</i> sp. <i>Paenibacillus kribbensis</i>	Organic acids.	(Marra et al., 2015)
Haloarchaea	<i>Haloarcula argentinensis</i> ; <i>Halobacterium</i> sp.; <i>Halococcus</i> sp.; <i>Halococcus hamelinensis</i> ; <i>Haloferax</i> sp., <i>H.</i> <i>alexandrines</i> ; <i>H. larsenii</i> ; <i>H. volcanii</i> <i>Halolamina</i> sp.; <i>Halolamina pelagic</i> ; <i>Halosarcina</i> sp.; <i>Halostagnicola</i> <i>kamekura</i> .; <i>Haloterrigena</i> <i>sp.</i>	Organic acids.	(Yadav et al., 2015)
Fungi	<i>Aspergillus niger</i> ; <i>Penicillium pinophilum</i> ; <i>Talaromyces rotundus</i> ; <i>Penicillium pimiteouiense</i> ; <i>Acremonium strictum</i> ;	Organic acids.	(Chuang et al., 2006) (Klaic et al., 2017)
Fungi	<i>Penicillium chrysogenum</i> ; <i>Aspergillus awamori</i>	Organic acids. Phosphatase and phytase enzymes.	(Babana et al., 2013)
Fungi	<i>Aspergillus niger</i> ; <i>Aspergillus flavus</i> ; <i>Trichoderma harzianum</i>	Enzymes: phytase, acid phosphatase + phytase; carboxymethyl cellulase + phosphatase.	(Gaiind, 2014)
Fungi	<i>Paecilomyces lilacinus</i>	Organic acids.	(Hernández-Leal et al., 2011)
Fungi	<i>Aspergillus niger</i> ; <i>Penicillium</i> <i>brevicompactum</i>	Enzymes: phytases. Organic acids.	(Perea et al., 2019)
AMF	<i>Acaulospora scrobiculata</i> ;; <i>Glomus</i> sp.; <i>G. deserticola</i> ; <i>G. intraradices</i> ; <i>G. versiforme</i>	Enhances the uptake of soluble phosphates in mycorrhizal association.	(Maaloum et al., 2020)
AMF	<i>Rhizophagus intraradices</i>	Enhances the uptake of soluble phosphates in mycorrhizal association.	(Velzquez et al., 2017)
AMF	<i>Acaulospora morrowiae</i> ; <i>Acualospora spinose</i> ; <i>Acualospora scrobiculata</i> ; <i>Gigaspora rosea</i> ; <i>Scutellospora pellucida</i> ; <i>Glomus macrocarpum</i> ; <i>Funnelformis mosseae</i> ; <i>Funnelformes geosporum</i> ; <i>Rhizophagus aggregatus</i> ;	Enhances the uptake of soluble phosphates in mycorrhizal association.	(Perea et al., 2019)

**Table 3.** Total P and bioavailable P values in different types of compost and vermicompost

Composts	Total phosphorus (TP) (g/kg)	Bioavailable phosphorus (PAV) (g/kg)	Reference
Urban solid organic waste	1.99	ND	(García-Molano et al., 2021)
Wastewater sludge	6.00	2.51	(Grigatti et al., 2017)
Rice straw	8.50	0.90	
Wheat straw	7.80	0.80	
Mustard stubble	6.10	0.75	(Moharana et al., 2020)
Chickpea stubble	6.50	0.70	
Tree branches	6.30	0.62	
Poultry litter without inoculation	230.00	9.69	
Poultry litter with <i>Pseudomonas sp.</i>	410.00	17.20	(Billah and Bano, 2014)
Poultry litter with <i>Proteus sp.</i>	300.00	12.40	
Pig manure	21.95	6.08	
Chicken manure	29.99	8.64	
Municipal solid waste	6.76	1.17	(Wei et al., 2015)
Kitchen waste	3.36	0.78	
Green waste	4.62	1.86	
Rice straw with pig manure	1.34	ND	(Chen and Wan, 2023)
Municipal solid waste	3.00	ND	(Ahmadi et al., 2020)
Organic waste	2.10	ND	(Sandoval et al., 2020)
Organic fraction of urban solid waste	5.40	ND	(Saldarriaga et al., 2018)
Agro-industrial waste	3.20-8.90	ND	(Hernández-Lara et al., 2021)
Kitchen waste	4.58	0.59	(Zhan et al., 2021)
Ruminal content	15.10	ND	(Bohórquez-Sandoval et al., 2020)
Grape pomace waste:			
Almariño variety	0.75	ND	(Gómez-Brandón et al., 2021)
Mencía variety	1.41		
Mixture of weed waste, manure, and sawdust	12.87	ND	(Kausar and Khwairakpam, 2022)
Agro-industrial poultry waste	16.90	ND	(Niedzialkoski et al., 2021)
Vineyard waste	2.00	ND	(Blaya et al., 2013)
Inoculated residual municipal solid waste with PSB	10-18	4.20-8.10	(Wei et al., 2017)
Rice straw and PR	24.20	1.15	
Wheat straw and PR	22.70	1.08	
Mustard stubble and PR	21.00	1.08	(Moharana et al., 2020)
Chickpea stubble and PR	21.60	1.08	
Tree branches and PR	20.90	0.95	
Kitchen waste with PR 5%	20.63	0.29	
PR 10%	31.16	0.74	(Zhan et al., 2021)
PR 15%	40.64	0.45	
Mixture of manures: Cattle, horse, and poultry, adding PR in three concentrations:			
Compost with 5% PR	145.00	1.60	
Compost with 10% PR	151.00	1.65	(Páez et al., 2022)
Compost with 15% PR	215.00	1.70	
Vineyard waste:			
With <i>Trichoderma harzianum</i> inoculation	1.50	ND	(Blaya et al., 2013)
Plant and animal waste with <i>Azospirillum brasilense</i> inoculation	ND	12.00	(Hafez et al., 2021)
Rice straw + cattle manure	ND	0.62	
Rice straw + farmyard manure	ND	0.74	
Rice straw + poultry manure	ND	1.48	
Rice straw + cattle manure + PSF	7.12	0.85	(Gaid, 2014)
Rice straw + farmyard manure + PSF	6.70	0.85	
Rice straw + poultry manure + PSF	9.98	1.56	

PR: Phosphate rock; PSF: Phosphate-solubilizing fungi; PSB: Phosphate-solubilizing bacteria.

2021; Velzquez et al., 2017). The phosphorus absorbed by plants is mineralized organic phosphorus, which originates from phosphate esters where P is bound to oxygen in its maximum oxidation state (+5), making it highly suscepti-

ble to mineralization. There are also phosphonates, where phosphorus is directly bound to carbon, forming bonds that require more energy to break, operating with an oxidation state of +3 (Tapia-Torres and García-Oliva, 2013).

### Quantification methods of Phosphorous forms

Various methodologies for extracting plant-available P have been developed, among which the most relevant are Colwell (1963), Bray I and II (1945), Mehlich (1984), and Olsen-P (1954); the latter being the most widely used analytical method (Milham et al., 2023). Moharana et al. (2020) indicate that this method is commonly used due to its response to different substrates, making it a good indicator of plant-available P; In some cases, another indicator of availability is citric acid-soluble P at 2% for acidic media (CASP); it is also important to determine the percentage of P solubilization. In response to this, the European Commission proposed the SMT (Standards, Measurements, and Testing) method, which analyzes total phosphorus (TP), inorganic phosphorus (Pi), organic phosphorus (Po), non-apatite phosphorus (NAP), and apatite phosphorus (AP), aiming for uniformity in fractionation methods (Ruban et al., 2001; García, 2014; Velasco, 2021).

### Phosphorus availability in organic waste composts

The content and availability of phosphorus in composts can increase during the composting process, as indicated by Cheng and Wan (2022), 2023. For instance, compost made from sorghum straw and pig manure exhibited an increase from an initial phosphorus content of 0.5 g/kg to 1.34 g/kg after composting (Table 3). The phosphorus content in composts is influenced by the raw materials used, suggesting that combining various materials can enhance the macronutrient values (Table 3). This results in increased phosphorus content and promotes microorganisms involved in solubilization or mineralization, facilitating the gradual release of phosphorus and preventing losses through leaching or chelation (Vicentin et al., 2021).

Ortega (2022) demonstrated the production of enzymes like phytases, acid phosphatases, alkaline phosphatases, and neutral phosphatases by inoculating *Pseudomonas aeruginosa* into cattle manure compost. This led to the release of orthophosphate ions, highlighting the synthesis of various enzymes with mineralizing functions. Wan et al. (2021) showed that alkaline phosphatase activity was intense toward the end of composting, potentially favoring increased available phosphorus from organic waste. However, other enzymatic activities, such as  $\beta$ -1,4-glucosidase,  $\beta$ -D-fructofuranosidase,  $\beta$ -1,4-N-acetylglucosaminidase, and sulfatase, may also influence phosphorus availability, although this relationship has not been extensively studied. Wei et al. (2018b) and Wei et al. (2018a) found that formic acid exhibited the most solubilization activity for various forms of phosphorus, including TP, Po, Pi, Olsen-P, PAC (phosphorus soluble in citric acid), and microbial biomass phosphorus (MBP). Oxalic acid acted on TP, Po, Pi, and MBP, while citric acid influenced Po and MBP. These activities occurred during the cooling phase of composting, suggesting that the Organic acids, available phosphorus, and phosphate-solubilizing microorganisms reached the soil, enhancing plant activity.

The total phosphorus content in composts varies depending on the source material, ranging from 2 to 9 g/kg for composts derived from organic waste, sludge, and agro-

industrial wastes, to 14 to 17 g/kg for composts produced from manures or animal wastes (Table 3). Post-composting, phosphorus tends to accumulate, with substrate influencing microbial community establishment, particularly in animal manure composts, where phosphorus levels can reach 34 kg/ton (Williams, 2013). However, the majority of available phosphorus comes from organic matter, with content varying based on phosphorus form, carbon, and nitrogen levels (Hernández-Leal et al., 2011). Understanding the native microbiota of composted materials is essential for enhancing process efficiency (Gaiind, 2014). Wei et al. (2016) identified key phylogenetic groups involved in phosphorus transformation, such as *Firmicutes*, *Proteobacteria*, and *Bacteroidetes*, which contribute to the formation of slow-release phosphorus sources for plants. Families like *Pseudomonadaceae*, *Enterobacteriaceae*, and *Bacillaceae* are capable of solubilizing phosphorus in soils and are commonly found in composts and plant rhizospheres. Greater raw material diversity corresponds to increased microbial diversity and enzyme production. When compost is applied to the soil, it promotes mineralization processes of organic phosphorus in the soil's organic matter, laying the foundation for developing techniques to isolate microorganisms synthesizing phosphatases for use as biofertilizers.

The total phosphorus content or availability in composts inoculated with phosphate-solubilizing microorganisms (PSMs) varies based on the raw material type and specific PSMs used. Billah and Bano (2014) observed superior outcomes with the phosphate-solubilizing bacterium *Pseudomonas sp.*, compared to *Proteus sp.* and the control without inoculation, resulting in increased plant height and grain production. Biochar + oil palm bunch compost applied to maize crops fosters populations of phosphate-solubilizing fungi like *Aspergillus sp.* and *Neosartorya sp.*, enhancing microorganism adaptability and promoting maize growth through increased phosphorus absorption. This medium increased phosphorus absorption in maize plants by 100 – 200%, promoting their growth by producing various compounds. Chen and Wan (2023) highlighted the challenges of decomposing residues with high lecithin levels and demonstrated the mineralization of organic phosphorus (Po) by the WWJ-22 strain of *Pseudomonas sp.*, leading to increased phosphorus availability. Yadav et al. (2017) demonstrated that compost from various sources, including agricultural wastes and manure, with added phosphate rock (PR) and inoculated heat-resistant phosphate-solubilizing bacteria (*Brevibacillus* and *Bacillus*) in wheat cultivation, resulted in elevated total phosphorus levels, shoot and root length, and plant biomass. They emphasized compost as a conditioning medium for phosphate-solubilizing bacteria (PSB), essential for slowly solubilizing phosphorus for plant growth. Babana et al. (2013) compared the solubilization of Tilemsi phosphate rock using bacteria (*Pseudomonas sp.* and *Vibrio splendidus*) and fungi (*Penicillium chrysogenum thom* and *Agrobacterium tumefaciens*) isolated from wheat rhizospheres. Bacteria showed a higher solubility index, but fungi had more elevated solubilized phosphorus (Sarr et al., 2020). The latter demonstrated that heat-resistant phosphate-solubilizing fungi, by the end of composting,

had a stronger correlation with labile phosphorus (labile-P) and total phosphorus (TP) than bacteria, along with a higher correlation with alkaline phosphatase activity.

Gaïnd (2014) observed the concentration of mineralized phosphorus (Po), as measured by Olsen-P, doubling the phosphorus compared to the control treatment in different types of composts (rice straw with poultry manure, rice straw with cattle manure, and rice straw with farmyard manure) inoculated with a consortium of fungi (*Aspergillus niger*, *Aspergillus flavus*, and *Trichoderma harzianum*). In another study, Hafez et al. (2021) inoculated the bacterium *Azospirillum brasilense* into organic ecological waste compost, resulting in a notable increase in labile phosphorus, highlighting the role of indoleacetic acid (IAA) metabolic pathways in facilitating phosphorus solubilization and enhancing crop production (Table 3).

Additionally, vermicomposting of organic waste contributes phosphorus (P) as an organic fertilizer, facilitated by the mineralization of organic phosphorus within the earthworm's digestive tract and the production of enzymes like phosphatases and phytases. Studies by Bohórquez-Sandoval et al. (2020) and Gómez-Brandón et al. (2021) demonstrated increased total phosphorus (TP) content in vermicomposts derived from different organic materials. Combining composting with vermicomposting has also been explored, revealing promising phosphorus levels in the mix (Kausar and Khwairakpam, 2022; Niedzialkoski et al., 2021; Zuhair et al., 2022).

Related to the P absorbed by plants attending to its form, Moreno and Moral (2007) showed that only 20% to 40% of the total phosphorus (P) content is plant-available. This fraction represents labile phosphorus, which constitutes part of the bioavailable phosphorus fractions for plants. The dynamics in soils and, consequently, in crops depend on these contributions. Similarly, phosphorus levels play a crucial role in establishing the microbiota since phosphorus is present in cellular organelles or forms a part of ATP, the primary molecule for cellular energy acquisition. Consequently, plants can also mineralize phosphorus for their metabolic functions (Moreno and Moral, 2007).

Yang et al. (2018) found that cattle manure and maize stalk compost applied to soybean cultivation increased the density of arbuscular mycorrhizal fungi (AMF) spores and hyphae. This enhancement favored the flowering and maturation phases of the crop, regardless of the quantity of compost added. These findings suggest that the fungal spores belong to the native soil population of the *Glomeraceae* family. They highlighted that root colonization is influenced by the soybean crop's demand for substantial amounts of phosphorus (P) for atmospheric nitrogen fixation. Furthermore, the hyphae promote a slow and sustained release of labile P for the plant.

### 3. Conclusion

The need to find alternative sources of P to maintain sustainable agriculture has turned organic waste composts and sewage sludge into a nutrient source to reduce the use of inorganic sources. Depending on the raw materials used, and also due to the incorporation of phosphorus-solubilizing

microorganisms (PSM), their contents can increase as these microorganisms are involved in their solubilization. There are different mechanisms of P solubilization, such as Organic acids, phosphatases, symbiotic relationships, microbial interactions, and the action of phosphorus-related genes, which increase its slow release, reducing the use of highly soluble sources.

#### Author contribution

The authors confirm the conception and design of the study: Bohórquez-Sandoval LJ, García-Molano JF, Pascual-Valero JA, and Ros-Muñoz M; data collection: Bohórquez-Sandoval LJ, García-Molano JF; analysis and interpretation of results: Bohórquez-Sandoval LJ, García-Molano JF, and Ros-Muñoz M; preparation of the preliminary manuscript: Bohórquez-Sandoval LJ, García-Molano JF. The results were evaluated by all authors, and the final version of the manuscript was approved. The authors and contribution to the research effort adhere to the authorship standards established in the IJROWA Authorship Guidelines and as advised by the Committee on Publication Ethics (COPE).

#### Availability of data and materials

The data that support the findings of this study are available from the corresponding author upon reasonable request.

#### Conflict of interest statement

The authors declare that they are no conflict of interest associated with this study.

#### Open access

This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the OICC Press publisher. To view a copy of this license, visit <https://creativecommons.org/licenses/by/4.0/>.

## References

- Adnan M, Shah Z, Fahad S, Arif M, Alam M, Khan IA, Mian IA, et al. (2017) Phosphate-solubilizing bacteria nullify the antagonistic effect of soil calcification on bioavailability of phosphorus in alkaline soils. *Sci Rep* 7 (1): 1–13. <https://doi.org/10.1038/s41598-017-16537-5>
- Aguado GA, Moreno B, Jiménez B, García E, Preciado RE (2012) Impacto de los sideróforos microbianos y fitosideróforos en la asimilación de hierro por las plantas: Una síntesis. *Rev Fitotec Mex* 35 (1): 9–21. <https://doi.org/10.35196/rfm.2012.1.9>
- Ahmadi T, Casas CA, Escobar N, García YE (2020) Municipal organic solid waste composting: Development of a tele-monitoring and automation control system. *Agron Res* 18 (3): 1911–25. <https://doi.org/10.15159/AR.20.212>
- Antoun H (2012) Beneficial microorganisms for the sustainable use of phosphates in agriculture. *Pro Eng* 46:62–67. <https://doi.org/10.1016/j.proeng.2012.09.446>
- Atoloye IA, Jacobson A, Creech E, Reeve J (2021) Variable impact of compost on phosphorus dynamics in organic dryland soils following a one-time application. *Soil Society of America J* 85 (4): 1122–38. <https://doi.org/10.1002/saj2.20275>
- Awasthi MK, Pandey AK, Bundela PS, Khan J (2015) Co-composting of organic fraction of municipal solid waste mixed with different bulking waste: Characterization of physicochemical parameters and microbial enzymatic dynamic. *Bioresour Technol* 182:200–207. <https://doi.org/10.1016/j.biortech.2015.01.104>
- Azam HM, Alam ST, Hasan M, Yameogo DDS, Kannan AD, Rahman A, Kwon MJ (2019) Phosphorous in the environment: characteristics with distribution and effects, removal mechanisms, treatment technologies, and factors affecting recovery as minerals in natural and engineered systems. *Environ Sci Pollut Res*, 2620183–20207. <https://doi.org/10.1007/s11356-019-04732-y>
- Babana AH, Dicko AH, Maïga K, Traoré D (2013) Characterization of rock phosphate-solubilizing microorganisms isolated from wheat (*Triticum aestivum* L.) rhizosphere in Mali. *J Microb Res* 1 (1): 1–6.
- Bank World (2018) Global waste to grow by 70 percent by 2050 unless urgent action is taken: World Bank Report Washington DC.
- Basílio F, Dias T, Santana MM, Melo J, Carvalho L, Correia P, Cruz C (2022) Multiple modes of action are needed to unlock soil phosphorus fractions unavailable for plants: The example of bacteria- and fungi-based biofertilizers. *Applied Soil Ecology* 178:104550. <https://doi.org/10.1016/j.apsoil.2022.104550>
- Behera BC, Yadav H, Singh SK, Sethi BK, Mishra RR, Kumari S, Thatoi H (2017) Alkaline phosphatase activity of a phosphate solubilizing *Alcaligenes faecalis*, isolated from Mangrove soil. *Biotech Res Inno* 1 (1): 101–111. <https://doi.org/10.1016/j.biori.2017.01.003>
- Beltrán-Pineda ME (2015) La solubilización de fosfatos como estrategia microbiana para promover el crecimiento vegetal. *Revis Cien y Tec* 15 (1): 101–113. [https://doi.org/10.21930/rcta.vol15\\_num1\\_art:401](https://doi.org/10.21930/rcta.vol15_num1_art:401)
- Billah M, Bano A (2014) Role of plant growth promoting rhizobacteria in modulating the efficiency of poultry litter composting with rock phosphate and its effect on growth and yield of wheat. *Waste Manag Res* 3 (1): 63–72. <https://doi.org/10.1177/0734242X14559593>
- Blaya J, López-Mondéjar R, Lloret E, Pascual JA, Ros M (2013) Changes induced by *Trichoderma harzianum* in suppressive compost controlling *Fusarium wilt*. 107 (1): 12–119. <https://doi.org/10.1016/j.pestbp.2013.06.001>
- Bohórquez-Sandoval L, García-Molano F, Murillo-Arango W, Cuervo-Bejarano J, Pulido-Soler N (2020) Vermicomposting: A transformation alternative for rumen content generated in slaughterhouses. *Rev Facul Nac Agron Medellín* 73 (2): 9201–12. <https://doi.org/10.15446/rfnam.v73n2.80104>
- Bolívar-Anillo HJ, Contreras-Zentella ML, Teherán-Sierra LG (2016) *Burkholderia tropica*: Una bacteria con gran potencial para su uso en la agricultura. 19 (2): 102–08. <https://doi.org/10.1016/j.recqb.2016.06.003>
- Bowen HJM (1979) Environmental chemistry of the elements. (p. 333pp). Dep. of Chem., Reading Univ., Reading RG6 2AH, UK. *Academic Press* ISBN 978-0-12-120450-1
- Castillo-Arteaga RD, Burbano-Rosero EM, Otero-Ramírez ID, Fernández-Izquierdo P (2016) Degradación de oxalato por bacterias oxalotróficas asociadas a plantas del género *Oxalis* sp en regiones Andinas del departamento de Nariño, Colombia. *Universidad y Salud sección de artículos originales Degradación* 18 (1): 69–78.
- Chen P, Wan W (2023) Rare alkaline phosphatase-harboring bacteria mediate organic phosphorus mineralization during swine manure composting. 368 (1): 128335. <https://doi.org/10.1016/j.biortech.2022.128335>
- Chen X, Yan X, Wang M, Cai Y, Weng X, Su D, Guo J, et al. (2022) Long-term excessive phosphorus fertilization alters soil phosphorus fractions in the acidic soil of pomelo orchards. *Soil Tillage Res* 215:105214. <https://doi.org/10.1016/j.still.2021.105214>
- Cheng Y, Wan W (2022) Alkaline phosphomonoesterase-harboring bacteria facilitate phosphorus availability during winter composting with different animal manures. *Journal Clean Prod* 376 (1): 134299. <https://doi.org/10.1016/j.jclepro.2022.134299>

- Chuang CC, Kuo YL, Chao CC, Chao WL (2006) Solubilization of inorganic phosphates and plant growth promotion by *Aspergillus Niger*. *Biol Ferti Soil* 43 (5): 575–84. <https://doi.org/10.1007/s00374-006-0140-3>
- Contrato CRUC (2018) Roca Fosfórica Caracterización y análisis de mercado internacional de minerales en el corto, mediano, y largo plazo con vigencia al año 2035 Accessed January 26:2022.
- Corbridge DE (2013) Phosphorus: chemistry, biochemistry and technology. *CRC Press LLC. Sixth edition. UK*
- Corrales LC, Arévalo ZY, Moreno VE (2014) Solubilización de fosfatos: Una función microbiana importante en el desarrollo vegetal. *Nova* 12 (21): 67. <https://doi.org/10.22490/24629448.997>
- Darch T, Blackwell M. SA, Chadwick D, Haygarth PM, Hawkins JMB, Turner BL (2016) Assessment of bioavailable organic phosphorus in tropical forest soils by organic acid extraction and phosphatase hydrolysis. *Geoderma* 284:93–102. <https://doi.org/10.1016/j.geoderma.2016.08.018>
- Djandja OS, Salami AA, Wang ZC, Duo J, Yin LX, Duan PG (2022) Random forest-based modeling for insights on phosphorus content in hydrochar produced from hydrothermal carbonization of sewage sludge. *Energy* 245:123295. <https://doi.org/10.1016/j.energy.2022.123295>
- Escamilla SR (2015) La historia del fósforo: Una reflexión acerca de la seguridad alimentaria mundial. *Boletín Cient Cien Bas e INg ICBI* 2, 4, <https://doi.org/10.29057/icbi.v2i4.548>
- Espinel NM (2020) Aprovechamiento de roca fosfórica, por vía térmica, para la obtención de termofosfatos. *Rev Inv* 12 (2): 113–133. <https://doi.org/10.29097/2011-639x.266>
- Estrada-Bonilla GA, Lopes CM, Durrer A, Alves PRL, Pasaglia N, Cardoso EJBN (2017) Effect of phosphate-solubilizing bacteria on phosphorus dynamics and the bacterial community during composting of sugarcane industry waste. *System Appl Microb* 40 (5): 308–13. <https://doi.org/10.1016/j.syapm.2017.05.003>
- Etesami H, Jeong BR, Glick BR (2021) Contribution of Arbuscular Mycorrhizal fungi, phosphate-solubilizing bacteria, and silicon to p uptake by plant. *Front Plant Sci* 12:1–29. <https://doi.org/10.3389/fpls.2021.699618>
- EU (2019) Regulation (EU) 2019/1009 of the European Parliament and of the Council of 5 June 2019. *Official Journal of the European Union*, 1–114.
- Ferenhof HA, Fernandes RF (2016) Desmistificando a revisão de literatura como base para redação científica: Método SSF Scientific Writing and Research Methods View project Organizational learning and unlearning View project Accessed December 29, 2021 <https://www.researchgate.net/publication/325070845>
- Fernández-Marcos ML (2011) Contaminación por fósforo procedente de la fertilización orgánica de suelos agrícolas. Gestión de residuos orgánicos de uso agrícola *Guía de residuos orgánicos de uso agrícola Santiago de Compostela Servizo de Publicacions Universidade de Santiago de Compostela p 25*
- Ferraz R, Almeida N, Andrade N de, Scheffer I, Tirloni B, Oliveira A de, Domínguez J, Josemar R (2022) Vermicomposting of cow manure: Effect of time on earthworm biomass and chemical, physical, and biological properties of vermicompost. *Bioresour Technol* 345:126572. <https://doi.org/10.1016/j.biortech.2021.126572>
- Fuller WH (1972) Phosphorus element and geochemistry. *The encyclopedia of geochemistry and environmental sciences: New York, Van Nostrand Reinhold Co*, 942–46.
- Gaünd S (2014) Effect of fungal consortium and animal manure amendments on phosphorus fractions of paddy-straw compost. *Int Bio Biodegrad* 94:90–97. <https://doi.org/10.1016/j.ibiod.2014.06.023>
- Galvis GJA (2016) Residuos sólidos: Problema, conceptos básicos y algunas estrategias de solución. *Revista Gestión y Región* 22:101–119.
- Gao X, Tan W, Zhao Y, Wu J, Sun Q, Qi H, Xie X, Wei Z (2019) Diversity in the mechanisms of humin formation during composting with different materials. *Environ Sci Technol* 53 (7): 3653–62. <https://doi.org/10.1021/acs.est.8b06401>
- García AM (2014) Residuos orgánicos como fuentes de fósforo (Tesis doctoral) pp. 35-36. Universidad Politécnica de Madrid
- García-Molano JF, Cuervo-Bejarano WJ, Rodolfi M, Jaramillo-Garda LS, Ganino T (2022) Can olive pruning forms influence the olive rhizosphere? The root microbiota and the rhizosphere properties in the alto ricaurte (Colombia). *Agronomy* 12 (5) <https://doi.org/10.3390/agronomy12051159>
- García-Molano JF, Parra-Alba J D, Guevara LAP (2021) Characterization of composted organic solid fertilizer and fermented liquid fertilizer produced from the urban organic solid waste in paipa, boyacá, colombia. *Int J Recycl Waste Org Agricul* 10 (4): 379–95. <https://doi.org/10.30486/ijrowa.2021.1901014.1083>
- Grigatti M, Boanini E, Biase G Di, Marzadori C, Ciavatta C (2017) Effect of iron sulphate on the phosphorus speciation from agro-industrial sludge based and sewage sludge based compost. *Waste Manag* 69:353–359. <https://doi.org/10.1016/j.wasman.2017.08.029>
- Gómez-Brandón M, Martínez-Cordeiro H, Domínguez J (2021) Changes in the nutrient dynamics and microbiological properties of grape marc in a continuous-feeding vermicomposting system. *Waste Manag* 69:353–359. <https://doi.org/10.1016/j.wasman.2017.08.029>

- Hafez M, El-Ezz SF Abo, Popov AI, Rashad M (2021) Organic amendments combined with plant growth-promoting rhizobacteria (*Azospirillum Brasilense*) as an eco-friendly by-product to remediate and enhance the fertility of saline sodic-soils in Egypt. 52 (12): 1416–14. <https://doi.org/10.1080/00103624.2021.1885687>
- Hernández-Berriel MC, Aguilar-Virgen Q, Taboada-González P, Lima-Morra R, Eljaiek-Urzola M, Márquez-Benavides L, Buenrostro-Delgado O (2016) Generación y composición de los residuos sólidos urbanos en América latina y el caribe. *Rev Int Contam Amb* 32 (1): 11–22. <https://doi.org/10.20937/RICA.2016.32.05.02>
- Hernández-Lara A, Ros M, Pérez-Murcia MD, Bustamante MA, Moral R, Andreu-Rodríguez FJ, Fernández JA, Egea-Gilabert C, Pascual JA (2021) The influence of feedstocks and additives in 23 added-value composts as a growing media component on *Pythium irregulare* suppressivity. *Waste Manag* 120:351–63. <https://doi.org/10.1016/j.wasman.2020.11.053>
- Hernández-Leal TI, Carrión G, Heredia G (2011) Solubilización in vitro de fosfatos por una cepa de *Paeclomyces lilacinus* (Thom) Samson. *Agrociencia* 45 (8): 881–892. <https://doi.org/10.1007/s10661-020-08437-5>
- Hoang AT, Varbanov PS, Nizetić S, Sirohi R, Pandey A, Luque R, Ng KH, Pham VV (2022) Perspective review on municipal solid waste-to-energy route: Characteristics, management strategy, and role in circular economy. *J Clean Prod* 359:131897. <https://doi.org/10.1016/j.jclepro.2022.131897>
- ICONTEC (2011) Norma Técnica Colombiana NTC-5167 *Productos orgánicos usados como abonos o fertilizantes y como enmiendas de suelo.*
- Iida CL, Shock CC (2011) El dilema del fósforo. O. S. U. *Panacea Catalog extensión Oregon State University* 16 (42): 189. [https://ir.library.oregonstate.edu/concern/open\\_educational\\_resources/b2773w01b](https://ir.library.oregonstate.edu/concern/open_educational_resources/b2773w01b)
- Kaur G (2020) Microbial phytases in plant minerals acquisition. In *Molecular Aspects of Plant Beneficial Microbes in Agriculture*. INC, <https://doi.org/10.1016/b978-0-12-818469-1.00016-x>
- Kauser H, Khwairakpam M (2022) Organic waste management by two-stage composting process to decrease the time required for vermicomposting. *ETI* 25:102193. <https://doi.org/10.1016/j.eti.2021.102193>
- Klaic R, Plotegher F, Ribeiro C, Zangirolami TC, Farinas CS (2017) A novel combined mechanical-biological approach to improve rock phosphate solubilization. *Int J Min Pro* 161:50–8. <https://doi.org/10.1016/j.minpro.2017.02.009>
- Kwesi-Asomaning S (2020) Processes and factors affecting phosphorus sorption in soils. *World J. Agric. Sci.* 45:1–16. <https://doi.org/10.5772/intechopen.90719>
- Licea-Herrera JI, Quiroz-Velasquez JDC, Hernández-Mendoza JL (2020) Impacto de *Azospirillum brasilense*, una rizobacteria que estimula la producción del ácido indol-3-acético como el mecanismo de mejora del crecimiento de las plantas en los cultivos agrícolas. *Rev Boliv Quim* 37 (1): 34–9. <https://doi.org/10.34098/2078-3949.37.1.5>
- Maaloum S El, Elabed A, Alaoui-Talibi Z, Meddich A El, Filali-Maltouf A, Douira A, Ibsouda-Koraichi S, Amir S, Modafar C El (2020) Effect of *Arbuscular Mycorrhizal* fungi and phosphate-solubilizing bacteria consortia associated with phospho-compost on phosphorus solubilization and growth of tomato seedlings (*Solanum lycopersicum* L.). *Communi Soil Sci Plan Analy* 51 (5): 622–34. <https://doi.org/10.1080/00103624.2020.1729376>
- Marra LM, Oliveira-Longatti SM de, Soares CRFS, Lima JM de, Olivares FL, Moreira FMS (2015) Initial pH of medium affects organic acids production but do not affect phosphate solubilization. *Brazil J Microb* 46 (2): 367–75. <https://doi.org/10.1590/S1517-838246246220131102>
- Milham PJ, Harvey N Carlson-Perret Morrison D, Andersson KO, Burkitt LL, Collins D, Haigh AM, Hannah MC, Tellinghuisen J, Dougherty WJ, Holford P (2023) *Social Sci Res Net World J. Agric. Sci.*, <https://doi.org/10.2139/ssrn.4461675>
- Moharana PC, Biswas DR, Ghosh A, Sarkar A (2020) Variability of crop residues determines solubilization and availability of phosphorus fractions during composting of rock phosphate enriched compost vis-à-vis ordinary compost. *Comm Soil Sci Plant Anal* 51 (15): 2085–01. <https://doi.org/10.1080/00103624.2020.1784921>
- Moreno CJ, Moral HR (2007) *Compostaje* (J. Moreno Casco, R. Moral Herrero, and E. Científicos.(eds.)). *Ediciones Mundi-prensa Libros ISBN* 8484764796
- Niedzialkoski RK, Marostica R, Damaceno FM, Costa LA, Costa MSS (2021) Combination of biological processes for agro-industrial poultry waste management: Effects on vermicomposting and anaerobic digestion. *Environ Manag* 297:113127. <https://doi.org/10.1016/j.jenvman.2021.113127>
- Nobile C, Lebrun M, Védère C, Honvault N, Aubertin ML, Faucon MP, Houben D (2022) Biochar and compost addition increases soil organic carbon content and substitutes P and K fertilizer in three French cropping systems. *Agro Sust Devel* 42 (6): 119. <https://doi.org/10.1007/s13593-022-00848-7>
- Oliveira V, Horta C, Dias-Ferreira C (2019) Evaluation of a phosphorus fertiliser produced from anaerobically digested organic fraction of municipal solid waste. *J Clean Pro* 238 <https://doi.org/10.1016/j.jclepro.2019.117911>

- Ordoñez YM, Fernandez BR, Lara LS, Rodríguez A, Uribe-Vélez D, Sanders IR (2016) Bacteria with phosphate solubilizing capacity alter mycorrhizal fungal growth both inside and outside the root and in the presence of native microbial communities. *PLOS ONE* 11 (6): 1–18. <https://doi.org/10.1371/journal.pone.0154438>
- Ortega AI (2020) Propuesta para el abordaje de las consecuencias en el ambiente del uso de compuestos del fósforo. Uso de compuestos del fósforo. *Rev Enseñ Quim* 3:170–188.
- Ortega TAE (2022) Adición de enzimas para la mineralización del fósforo contenido en los residuos agroindustriales orgánicos destinados al uso. *Disertación Universidad Autónoma de Queretaro*, <http://ri-ng.uaq.mx/handle/123456789/3514>
- Panhwar QA, Jusop S, Naher UA, Othman R, Razi MI (2013) Application of potential phosphate-solubilizing bacteria and organic acids on phosphate solubilization from phosphate rock in aerobic rice. *Sustain World J* 2103 (272409): 1–10. <https://doi.org/10.1155/2013/272409>
- Paredes-Mendoza M, Espinosa-Victoria D (2009) Organic acids produced by phosphate solubilizing rhizobacteria: A critical review. *Terra Latino Amer* 28:61–70.
- Pascual JA, García C, Hernandez T (1999) Comparison of fresh and composted organic waste in their efficacy for the improvement of arid soil quality. *Bioresour Technol* 68 (3): 255–264. [https://doi.org/10.1016/S0960-8524\(98\)00160-6](https://doi.org/10.1016/S0960-8524(98)00160-6)
- Pedrosa D, Farias TS, Pereira A, Tarcísio RA, Rêgo EF (2013) Monitoramento dos parâmetros físico-químicos na compostagem de resíduos agroindustriais monitoring of physical and chemical parameters in agroindustrial waste composting. *Pesquisas Agrárias e Ambientais* 1 (1): 44–48.
- Perea RY, Arias RM, Medel OR, Trejo AD, Heredia G, Rodríguez YY (2019) Effects of native Arbuscular mycorrhizal and phosphate-solubilizing fungi on coffee plants. *Agrofor Sys* 93 (3): 961–972. <https://doi.org/10.1007/s10457-018-0190-1>
- Páez LA, García JF, Parra JD, Lozano JL (2022) Effect of phosphoric rock on the chemical, microbiological and enzymatic quality of poultry, equine and cattle manure compost mix. *Int J Recycl Waste Org Agricul* 11:385–398. <https://doi.org/10.30486/IJROWA.2022.1930622.1247>
- Pérez-Cordero A, Tuberquia-Sierra A, Amell-Jímenez D (2014) Actividad in vitro de bacterias endófitas fijadoras de nitrógeno y solubilizadoras de fosfatos. *Agrono Mesoame* 25 (2): 213–223. <https://doi.org/10.15517/am.v25i2.15425>
- Pérez-Navarro O, Ley-Chong N, Rodríguez-Marroquí K, González-Suárez E (2016) Business opportunities of citric acid production by fermentation from sugar substrates in Cuba. *Revis Cen Azuc* 43 (2): 85–100.
- Pérez-Pazos JV, Sánchez-López DB (2017) Caracterización y efecto de Azotobacter, Azospirillum y Pseudomonas asociadas a Ipomoea batatas del Caribe Colombiano. *Rev Colom Biotecno* 19 (2): 35–46. <https://doi.org/10.15446/rev.colomb.biote.v19n2.69471>
- Rana R, Ganguly R, Gupta AK (2018) Physico-chemical characterization of municipal solid waste from Tricity region of Northern India: A case study. *J Mater Cycles Waste Manag* 20 (1): 678–89. <https://doi.org/10.1007/s10163-017-0615-3>
- Restrepo-Franco GM, Marulanda-Moreno S, Fe-Pérez Y De la, Osa A Díaz de la, Lucia-Baldani V, Hernández-Rodríguez A (2015) Bacterias solubilizadoras de fosfato y sus potencialidades de uso en la promoción del crecimiento de cultivos de importancia económica. *Revista CENIC. Cien Bioló* 41 (3): 185–88.
- Richardson K, Steffen W, Lucht W, Bendtsen J, Cornell SE, Donges JF, Rockström J (2023) Earth beyond six of nine planetary boundaries. *Sci Adv* 9 (37) <https://doi.org/10.1126/sciadv.adh2458>
- Ruban V, Lopez-Sánchez JF, Pardo P, Raurent G, Muntau H, Quevauviller PH (2001) Development of a harmonized phosphorus extraction procedure and certification of a sediment reference material. *J Environ Monitor* 3:121–25. <https://doi.org/10.1039/B005672N>
- Saeid A, Prochownik E, Dobrowolska-Iwanek J (2018) Phosphorus solubilization by *Bacillus species*. *Molecules* 23 (11): 1–18. <https://doi.org/10.3390/molecules23112897>
- Saldarriaga JF, Gallego JL, López JE (2018) Determination of kinetics parameters for composting process of the organic fraction of municipal solid waste separated at source. *Chem Eng Trans* 70:217–22. <https://doi.org/10.3303/CET1870037>
- Sandoval DA, Segura OJA, Rodríguez MJP (2020) Sustainable use of organic solid waste: Approach to a model developed by waste picker organizations. case study Bogotá DC Colombia. *Int J Engin Res Technol* 13 (8): 2067–2080. <https://doi.org/10.37624/ijert/13.8.2020.2067-2080>
- Santamaria S, Ferrera R (2002) Dinámica poblacional de *Eisenia andrei* (bouché 1972) en diferentes residuos orgánicos. *Terra Latinoamericana* 20 (3): 303–310.
- Santoro V, Schiavon M, Visentin I, Constán-Aguilar C, Cardinale F, Celi L (2021) Strigolactones affect phosphorus acquisition strategies in tomato plants. *Pl Cell Environ* 44 (11): 3628–3642. <https://doi.org/10.1111/pce.14169>
- Sarr PS, Tibiri EB, Fukuda M, Zongo AN, Compaore E, Nakamura S (2020) Phosphate-solubilizing fungi and alkaline phosphatase trigger the P solubilization during the co-composting of Sorghum straw residues with burkina faso phosphate rock. *Front Env Sci* 8:1–17. <https://doi.org/10.3389/fenvs.2020.559195>

- Silva LI, Pereira MC da, Carvalho AMX, Buttrós VH de, Pasqual M, J J Dória (2023) Phosphorus-solubilizing microorganisms: A key to sustainable agriculture. *Agriculture* 13 (2): 462. <https://doi.org/10.3390/agriculture13020462>
- Singh G, Goyne KW, Kabrick JM (2015) Determinants of total and available phosphorus in forested Alfisols and Ultisols of the Ozark Highlands, USA. *Geoderma Reg* 5:117–26. <https://doi.org/10.1016/j.geodrs.2015.05.001>
- Soobhany N (2018) Assessing the physicochemical properties and quality parameters during composting of different organic constituents of municipal solid waste. *J Environ Chem Eng* 6 (2): 1979–88. <https://doi.org/10.1016/j.jece.2018.02.049>
- Takahashi T, Dahlgren RA (2016) Nature, properties and function of aluminum–humus complexes in volcanic soils. *Geoderma* 263:110–21. <https://doi.org/10.1016/j.geoderma.2015.08.032>
- Tapia-Torres Y, García-Oliva F (2013) La disponibilidad del fósforo es producto de la actividad bacteriana en el suelo en ecosistemas oligotróficos: Una revisión crítica. *TERRA Latinoamericana*, 231–242.
- Tinoco-Varela D, Bayuelo-Jiménez JS (2021) Formas y distribución de fósforo en un Andisol con sistemas contrastantes de uso del suelo del centro de México. *Rev Terra Lat* 39:1–11. <https://doi.org/10.28940/terra.v39i0.881>
- Veith JA, Sposito G (1977) Reactions of Aluminosilicates, Aluminum Hydrous Oxides, and Aluminum Oxide with o-Phosphate: The formation of X-ray amorphous analogs of variscite and montebrasite. *Soil Sci Soc Am J* 41 (5): 870–76. <https://doi.org/10.2136/sssaj1977.03615995004100050011x>
- Velasco A (2021) Inoculación de compost con microorganismos solubilizadores de fosfato y su efecto sobre la disponibilidad del fósforo. 42
- Velzquez MS, Cabello MN, Elíades LA, Russo ML, Allegrucci N, Schalamuk S (2017) Combinación de hongos movilizadores y solubilizadores de fósforo con rocas fosfóricas y materiales volcánicos para la promoción del crecimiento de plantas de lechuga (*Lactuca sativa* L.). *Rev Arg Microb* 49 (4): 347–55. <https://doi.org/10.1016/j.ram.2016.07.005>
- Vera-Cardoso BC, Muñoz-Rojas J, Munive JA, Marín-Cevada V, Flores-Encarnación M, Carreño-López R (2017) Pirroloquinolinaquinona (PQQ) y las bacterias promotoras del crecimiento vegetal (PGPR). De la biosíntesis a los fenotipos. *Alianzas y Tendencias* 2 (1): 22–29. <https://doi.org/10.5281/zenodo.5081377>
- Vergara SE, Tchobanoglous G (2012) Municipal solid waste and the environment: A global perspective. *Ann Review Env Resour* 37:277–309. <https://doi.org/10.1146/annurev-environ-050511-122532>
- Vicentin R, Masín CE, Lescano MR, Zalazar CS (2021) Poultry litter stabilization by two-stage composting-vermicomposting process: Environmental, energetic and economic performance. *Chemosphere* 281:130872. <https://doi.org/10.1016/j.chemosphere.2021.130872>
- Wahid F, Fahad S, Danish S, Adnan M, Yue Z, Saud S, Siddiqui MH, Brtnicky M, Hammerschmidt T, Datta R (2020) Sustainable management with mycorrhizae and phosphate solubilizing bacteria for enhanced phosphorus uptake in calcareous soils. *Agriculture* 10 (8): 1–14. <https://doi.org/10.3390/agriculture10080334>
- Walton CR, Hao J, Huang F, Jenner FE, Williams H, Zerkle AL, Lipp AL, Hazen RM, Peters SE, Shorttle O (2023) Evolution of the crustal phosphorus reservoir. *Sci Adv* 9 (18) <https://doi.org/10.1126/sciadv.ade6923>
- Wan W, Wang Y, Tan J, Qin Y, Zuo W, Wu H, He H, He D (2021) Alkaline phosphatase-harboring bacterial community and multiple enzyme activity contribute to phosphorus transformation during vegetable waste and chicken manure composting. *Bioresour Technol* 297:122406. <https://doi.org/10.1111/pce.14169>
- Wei Y, Wei Z, Cao Z, Zhao Y, Zhao X, Lu Q, Wang X, Zhang X (2016) A regulating method for the distribution of phosphorus fractions based on environmental parameters related to the key phosphate-solubilizing bacteria during composting. *Bioresour Technol* 211:610–17. <https://doi.org/10.1016/j.biortech.2016.03.141>
- Wei Y, Zhao Y, Fan Y, Lu Q, Li M, Wei Q, Zhao Y, Cao Z, Wei Z (2017) Impact of phosphate-solubilizing bacteria inoculation methods on phosphorus transformation and long-term utilization in composting. *Bioresour Technol* 241:134–41. <https://doi.org/10.1016/j.biortech.2017.05.099>
- Wei Y, Zhao Y, Lu Q, Cao Z, Wei Z (2018a) Organophosphorus-degrading bacterial community during composting from different sources and their roles in phosphorus transformation. *Bioresour Technol* 264:277–84. <https://doi.org/10.1016/j.biortech.2018.05.088>
- Wei Y, Zhao Y, Shi M, Cao Z, Lu Q, Yang T, Fan Y, Wei Z (2018b) Effect of organic acids production and bacterial community on the possible mechanism of phosphorus solubilization during composting with enriched phosphate-solubilizing bacteria inoculation. *Bioresour Technol* 247:190–99. <https://doi.org/10.1016/j.biortech.2017.09.092>
- Wei Y, Zhao Y, Xi B, Wei Z, Li X, Cao Z (2015) Changes in phosphorus fractions during organic wastes composting from different sources. *Bioresour Technol* 189:349–56. <https://doi.org/10.1016/j.biortech.2015.04.031>

- Williams CM (2013) Características de la gallinaza de las aves de corral. North Carolina State University, Department of Poultry Science, Raleigh, NC, Estados Unidos de América. Función de las aves de corral en la nutrición humana, p.59. Gestión de residuos de aves de corral en los países en desarrollo, p. 53. *Revisión del desarrollo Agrícola (FAO)*
- Xu S, Jia K, Zheng Y, Chen W, Wang Z, Wei D, Sun B, et al. (2023) Phosphorus transformation behavior and phosphorus cycling genes expression in food waste composting with hydroxyapatite enhanced by phosphate-solubilizing bacteria. *Bioresour Technol* 376:128882. <https://doi.org/10.1016/j.biortech.2023.128882>
- Yadav AN, Sharma D, Gulati S, Singh S, Dey R, Pal KK, Kaushik R, Saxena Ak (2015) Haloarchaea endowed with phosphorus solubilization attribute implicated in phosphorus cycle. *Sci Report* 5:1–10. <https://doi.org/10.1038/srep12293>
- Yadav H, Fatima R, Sharma A, Mathur S (2017) Enhancement of applicability of rock phosphate in alkaline soils by organic compost. *Apply Soil Eco* 113:80–85. <https://doi.org/10.1016/j.apsoil.2017.02.004>
- Yang W, Gu S, Xin Y, Bello A, Sun W, Xu X (2018) Compost addition enhanced hyphal growth and sporulation of Arbuscular mycorrhizal fungi without affecting their community composition in the soil. *Front Microbiol* 9:1–13. <https://doi.org/10.3389/fmicb.2018.00169>
- Zhan Y, Zhang Z, Ma T, Zhang X, Wang R, Liu Y, Sun B, et al. (2021) Phosphorus excess changes rock phosphate solubilization level and bacterial community mediating phosphorus fractions mobilization during composting. *Bioresour Technol* 337:125433. <https://doi.org/10.1016/j.biortech.2021.125433>
- Zhang R, Gu J, Wang X, Li Y, Zhang K, Yin Y, Zhang X (2018) Contributions of the microbial community and environmental variables to antibiotic resistance genes during co-composting with swine manure and cotton stalks. *J Hazard Mater* 358:82–91. <https://doi.org/10.1016/j.jhazmat.2018.06.052>
- Zhang T, Wu X, Shaheen SM, Rinklebe J, Bolan NS, Ali EF, Li G, Tsang DCW (2021) Effects of microorganism-mediated inoculants on humification processes and phosphorus dynamics during the aerobic composting of swine manure. *J Hazard Mater* 416:0–2. <https://doi.org/10.1016/j.jhazmat.2021.125738>
- Zuhair RM, Assiuty ANI I, Khalil MA, Salama WM (2022) Efficacy of vermicompost amended and bacterial diversity on plant growth and pathogen control. *Int J Recycl Waste Org Agricul* 11 (1): 131–141. <https://doi.org/10.30486/ijrowa.2021.1919606.1176>