

Accepted manuscript (author version)

To appear in: International Journal of Recycling of Organic Waste in Agriculture (IJROWA)

Online ISSN: 2251-7715

Print ISSN: 2195-3228

This PDF file is not the final version of the record. This version will undergo further copyediting, typesetting, and review before being published in its definitive form. We are sharing this version to provide early access to the article. Please be aware that errors that could impact the content may be identified during the production process, and all legal disclaimers applicable to the journal remain valid.

Received: 14 Nov 2023

Revised: 06 Feb 2024

Accepted: 04 Apr 2024

DOI: 10.57647/ijrowa-m1j8-w486

Review

Systematic Literature Review on the Utilization of Tuber Crop Skins in the Context of Circular Agriculture

Aziz Caliskan¹, Norhidayah Abdullah^{1*}, Noriza Ishak¹, Divine Senanu Ametefe², Imene Tatar Caliskan³

¹ Faculty of Hotel and Tourism Management, Universiti Teknologi MARA (UiTM), 40450, Shah Alam, Malaysia.

² College of Engineering, School of Electrical Engineering, Universiti Teknologi MARA (UiTM), 40450 Shah Alam, Selangor, Malaysia.

³ Institute for advanced studies, Centre for Research in Biotechnology for Agriculture (CEBAR), Universiti Malaya, 50603, Kuala Lumpur, Malaysia.

*Corresponding author email: norhi813@uitm.edu.my

Abstract

Purpose: This Systematic Literature Review (SLR) investigates the utilization of tuber crop skins, specifically from yam, potato, taro, and cassava as an underexplored resource in circular agriculture. The study aims to assess their multifaceted applications across sectors such as animal feed, bioenergy, material science, wastewater treatment, and food innovation, addressing the overarching research question of how tuber crop skins can contribute to a more sustainable agricultural model.

Method: A rigorous review protocol was applied to analyze 26 articles sourced from ScienceDirect and Web of Science. The selection criteria focused on works that explore the application of tuber crop skins in various domains, ensuring a comprehensive coverage of the subject matter.

Results: The analysis reveals that tuber crop skins offer significant benefits, including serving as nutrient-rich livestock feed, potential bioenergy feedstocks, and a basis for bio-based plastics and biocomposites due to their high cellulose, hemicellulose, and lignin content. Their capacity for absorbing pollutants also highlights their utility in wastewater treatment. Additionally, their application in food fortification and the creation of functional foods showcases their versatility.

Conclusion: Despite the promising applications of tuber crop skins, there is a notable lack of comprehensive studies examining their long-term sustainability, scalability, and socio-economic impacts. The research identifies a crucial need for interdisciplinary studies to address these gaps, advocating for further exploration to harness the full potential of tuber crop skins in promoting sustainable circular agricultural practices.

Keywords: Agro-waste; Agricultural residues; Agricultural by-product; Sustainable resource management; Waste management; Nutrient recovery

Introduction

Sustainable agriculture has increasingly ascended to the forefront of international dialogue, commanding attention from policymakers, researchers, and stakeholders across multiple sectors. The urgency of this discourse is propelled by a confluence of factors, such as; escalating global population, diminishing natural resources, and the exigencies of climate change. These factors all converging to demand a radical rethinking of how we manage our food systems (Muhammad et al. 2019). Within this system, tuber crops such as potatoes, yams, cassava and taro occupy a critical aspect. These crops are more than mere dietary staples in numerous cultures; they are pivotal contributors to caloric and nutritional sustenance, effectively acting as essentials in global food security (Scott 2021; Zierer et al. 2021). However, amid the extensive research landscapes addressing cultivation practices (Dahal et al. 2019), disease resistance (Lastochkina et al. 2020; Lal et al. 2021), and nutritional profiles of these crops (Rinaldo 2020), there exists a somewhat neglected domain: the sizable waste generated from tuber skins. A considerable proportion of these skins are discarded or underutilized, often being relegated to low-value applications such as animal feed or, worse, landfill deposits (Sampaio et al. 2020; Wang et al. 2020a; Kayiwa et al. 2021; El-Sawi et al. 2023). This current model of disposal or limited use is economically suboptimal and ecologically unsustainable. It amplifies waste management challenges and intensifies the environmental footprint of agricultural operations. Even more disconcerting is that this dismissal of tuber crop skins overlooks their latent value. Preliminary studies have indicated that these skins are endowed with valuable nutrients and bioactive compounds such as antioxidants and phenols (Wang et al. 2020b). These elements could find applications in diverse sectors, ranging from food fortification to pharmaceuticals and even cosmeceuticals. This transformation converts what is currently considered 'waste' into a high-value commodity.

The indiscriminate disposal of tuber crop skins constitutes not merely a loss of potential value-addition but also worsens the environmental load by contributing to environmental degradation (El-Sawi et al. 2023). This directly contradicts the objective of circular agriculture, which advocate for a zero-waste, involving a closed-loop system that optimally utilizes all agricultural outputs (Sarwono 2023). In this sort of paradigm, waste from one process becomes input for another, thereby maximizing resource efficiency, reducing environmental degradation, and fostering economic sustainability (Cainelli et al. 2020). Despite its conceptual appeal and practical merits, circular agriculture as applied to tuber crop skins remains an underexplored area of inquiry. This limited

exploration creates a research gap that is both timely and significant. Addressing this gap does not just contribute to theoretical understandings of sustainable agriculture; it has profound implications for practice and policy, offering a multifaceted solution to challenges of waste management, resource utilization, and environmental conservation.

Consequently, the overarching objective of this study is to present a comprehensive evaluation of the prospects and modalities for integrating tuber crop skins into circular agricultural systems. By doing so, this study aims to highlight both the untapped potential and the practical methodologies for leveraging these crop by products. It views them as valuable resources rather than waste. This approach contributes a novel dimension to the domain of sustainable agriculture literature.

Scope of the Study

The scope of the study is defined to offer both a comprehensive and focused examination of the utility of tuber crop skins in the context of circular agriculture, concentrating on four principal tuber crops: potatoes (*Solanum tuberosum*), yams (*Dioscorea spp.*), cassava (*Manihot esculenta*), and taro (*Colocasia esculenta*). These crops were specifically chosen for their global ubiquity and significant contribution to food systems. Their variances in biochemical composition make them robust case studies for a diversified yet targeted analysis. Although other tubers exist, the focus on these four aims to lay a foundational understanding that could be inferred or adapted for other tuber varieties in future scholarly endeavours. In terms of geography, this review adopts a global perspective, by not placing emphasis on only regions where these crops are either staple foods or hold substantial economic significance.

Objective of the study

The primary goal of this study is to comprehensively evaluate the current state of research related to the utilization of tuber crop skins in the context of circular agriculture. To accomplish this, the study is guided by the following specific objectives:

- **To identify existing research on the utilization of tuber crop skins:** The first objective seeks to compile and assess the existing body of literature that discusses the various uses of tuber crop skins. This includes but is not limited to their nutritional profile, bioactive components, and current applications in food, pharmaceuticals, or other industries.
- **To assess the feasibility and efficiency of integrating tuber crop skins into circular agriculture practices:** This objective aims to evaluate how these uses align with the principles of circular agriculture. Special attention will be given to studies or data that specifically explore three key areas: the reduction of waste, optimization of resource use, and enhancement of economic benefits. This focus applies when integrating tuber crop skins into circular models.
- **To highlight gaps and future research needs:** Finally, this objective is geared towards identifying areas where existing research is lacking, inconclusive, or contradictory. Through a systematic analysis of the literature, this review will highlight gaps in current knowledge and propose avenues for future scholarly work.

Motivation of the study

The need for scholarly inquiry into waste minimization and value augmentation in tuber crops is motivated by multiple intersecting trends. Foremost among these is the escalating concern over food waste as a key driver of environmental disorder, manifest in soil degradation, water resource misallocation, and greenhouse gas emissions (Pang et al. 2020; Redlingshöfer et al. 2020; Shalini and Raghavan 2021). This urgency is underscored by a rapidly increasing global population, which heightens the demand for food (Ojuederie et al. 2019; Mahapatra et al. 2022). At the same time, it intensifies the imperative for sustainability across the entire agricultural value chain. Tuber crops such as potatoes, yams, cassava, and taro, occupy a central role in this dialogue, given their substantial contributions to food systems in both developed and developing nations (Neela & Fanta 2019; Tadele 2019). Despite their significance, a clear research deficit exists regarding the waste of tuber skins. These discarded byproducts contain untapped economic and nutritional potential. Their rich profiles of nutrients and bioactive compounds are amenable to diverse applications, ranging from food fortification to pharmaceutical interventions (Elhassaneen et al. 2016; Asharuddin et al. 2017; Fernandez et al. 2020; Mulia et al. 2023). Against this backdrop, circular agriculture emerges as an innovative and pragmatic framework for transforming these challenges into opportunities by promoting a zero-waste, resource-efficient agricultural paradigm (Velasco et al. 2022). However, current literature is disproportionately skewed toward macro-level strategies in circular agriculture. This includes practices such as crop rotation and manure management, leaving an academic void specific to the utility of tuber crop skins (Sui et al. 2018; Moretti et al. 2020). As a result, this study seeks to fill this gap by focusing on the skins of selected key tuber crops. The aim is to elucidate hitherto underexplored avenues for waste mitigation and value optimization. Hence, offering empirical and theoretical substantiation to their integration into a more resilient and sustainable circular agricultural ecosystem.

Significance of the study

This study marks a significant stride in sustainable agriculture by redefining the utilization of tuber crop skins within a circular agricultural model. It challenges the conventional perception of these skins as mere waste, uncovering their overlooked potential. Focusing on globally significant tubers like potatoes, yams, cassava, and taro, the study showcases their complex biochemical profiles, revealing a wealth of untapped value. The detailed examination of their nutritional and bioactive properties opens a variety of applications beyond their current limited use. This bridges the gap between academic research and practical implementations in food fortification, pharmaceuticals, cosmeceuticals, among others. As a result, this aligns with the progressive trends in circular economy practices, positioning the study at the forefront of sustainable agricultural innovation. By emphasizing circular agriculture specifically in the context of tuber crop skins, the study contributes a novel perspective to the literature, advocating for sustainable, zero-waste agricultural methods. It identifies key gaps in existing research, setting the stage for future exploration, and offering new insights into waste reduction and value augmentation in tuber crops. Therefore, the implication of the study extends beyond academia, resonating with the urgent need for sustainable agricultural practices in the face of escalating global food demands. It proposes a transformative approach, suggesting a shift from traditional waste management to a model in which tuber crop skins are recognized as valuable resources. Thus, this study not only fills a critical academic gap but also serves as a drive for policy and practical shifts towards more sustainable, economically viable agricultural systems.

The remainder of this paper is divided into 5 sections. Section 2 delineates the methodological approach employed for the Systematic Literature Review (SLR). In Section 3, we delve into a comprehensive review and synthesis of the pertinent literature. Section 4 is dedicated to presenting and discussing the principal findings of the study. The limitations of the study are critically examined in Section 5. Finally, the paper culminates with conclusions in Section 6.

Methodology

To explore the underutilized potential and multifaceted applications of tuber crop skins within the context of circular agriculture, our inquiry is anchored by a Systematic Literature Review (SLR) methodology. This methodological choice is predicated on the SLR's proven efficacy in rigorously collating, scrutinizing, and synthesizing insights across a broad spectrum of scholarly literature. As a result, ensuring a comprehensive and unbiased assessment of the complex issues surrounding the utilization of tuber crop skins. Focusing on key crops such as potatoes, yam, cassava, and taro, our SLR aims to showcase the myriad potential of these often-discarded tuber crop skin within the confines of circular agriculture. By collating and dissecting this body of academic works, our SLR not only elevates the scholarly rigor but also serves as a crucial aspect in the evolving discourse on circular agricultural practices and resource-efficient food systems.

Article Selection Method

In our rigorous pursuit of exhaustive literature pertaining to the utilization of tuber crop skins in circular agriculture, we employed a dual-database search strategy, involving Web of Science (WoS) and ScienceDirect. Recognizing WoS's strength in multidisciplinary research coverage and ScienceDirect's reputation for its expansive collection of high-quality academic works (Abioye et al. 2021), this binary approach facilitated a comprehensive extraction of scholarly articles directly relevant to our research query. Operating within the stringent parameters set by PRISMA guidelines (Page et al. 2021), as showed in Fig. 1, we undertook the systematic review, ensuring both clarity and breadth in our findings. Given the dynamic landscape of circular agriculture and sustainable resource management, we focused our review on articles published between 2020 and 2023. This approach was to secure the most current perspectives and innovations concerning the utilization of skins from tuber crops like potatoes, yams, cassava, and taro. Also, recognizing that circular agriculture is an evolving field influenced by emerging technologies, policy shifts, and societal attitudes, this time frame guarantees that our study encapsulates the latest trends, challenges, and opportunities. Keywords steering our search included phrases such as "tuber crop skins," "yam skins," "cassava skin," "potato skins," "taro skins," "circular agriculture," "sustainable farming," "resource optimization," "waste reduction," and "value addition." These search terms were used in various combinations to retrieve the most relevant articles. Furthermore, to uphold the scholarly rigor of our review, stringent inclusion and exclusion criteria were established. Five specific inclusion metrics were defined to focus our review on articles that precisely align with our research objectives. Meanwhile, four exclusion metrics served to filter out studies that were either tangentially related or did not meet academic quality standards.

Inclusion and exclusion criteria

To ensure the highest relevance and academic rigor in our study focusing on the utilization of tuber crop skins in circular agriculture, we meticulously crafted our selection parameters. Each chosen article had to meet all the inclusion criteria while avoiding any features that would place it under the exclusion benchmarks.




Figure 1. Flowchart of the systematic review process.

This rigorous approach ensures that the selected studies precisely illuminate the potential utilization of tuber crop skins. It focuses specifically on potatoes, yams, cassava, and taro within the circular agricultural sphere. Through this, we aim to fill the existing gaps in the literature, offering insights into how these often-overlooked tuber crop skins can contribute to both waste reduction and value addition in sustainable agriculture. Table 1 highlights the various predefined inclusion and exclusion criteria.

Table 1. Summary of the inclusion and exclusion criteria.

Inclusion Criteria		Exclusion Criteria	
Consideration	Description	Consideration	Description
Relevance to Research Objectives	Articles must focus on the utilization of tuber crop skins in the context of agriculture, specifically within the scope of circular or sustainable models.	Irrelevance to Objectives	Articles not focusing on the skin of tuber crops or circular agriculture will be excluded
Peer-Reviewed Publications	Only articles published in peer-reviewed journals will be considered, as they generally meet higher quality standards	Non-Peer-Reviewed Materials	Grey literature, opinion pieces, and non-peer-reviewed publications will be excluded
Language	Articles published in English will be included to ensure accurate interpretation and analysis	Non-English Publications	Articles published in languages other than English will be excluded due to potential interpretation challenges
Methodological Rigor	Studies must exhibit methodological rigor, including clear research designs, methodologies, and evidence-based conclusions	Out of Scope	Articles that are not focused on the any of the four tuber crop skins (i.e., yam, cassava, potato & taro)
Interdisciplinary Perspectives	Articles from a range of disciplines including agriculture, environmental science, and sustainability will be considered to provide a multifaceted perspective		

By explicitly stating these criteria, the study aims to eliminate biasness in article selection and ensure that only high-quality and relevant research is included in the review. This enhances the credibility and reliability of the findings and offers a pathway for potential replication and extension of the study.

Selected articles

Upon applying our carefully selected keywords and phrases, we initially gathered 143 publications: 97 from ScienceDirect and 46 from WoS, pertaining to the utilization of tuber crop skins in circular agriculture. After eliminating 26 duplicates, 117 articles remained for rigorous scrutiny. Our first-level screening involved a thorough examination of titles and abstracts. This led to the exclusion of 74 articles that did not directly address the opportunities and challenges of incorporating tuber crop skins. This specifically refers to the skins of potatoes, yams, cassava, and taro into circular agricultural systems. Upon delving into the full texts of the remaining 43 articles, 18 failed to meet our stringent inclusion criteria, while 1 was inaccessible. This systematic selection process cut our number of studies down to 24 relevant articles. To ensure exhaustive coverage, we utilized both backward and forward search techniques (Wolfswinkel et al. 2017; John et al. 2023), adding another 2 relevant articles to our collection. Consequently, our SLR consists of a curated selection of 26 journal articles that clearly align with our rigorous inclusion and exclusion benchmarks.

Review on the utilization of tuber crop skins in circular agriculture

Utilization of tuber crop skins has surfaced as an innovative avenue within the rapidly growing field of circular agriculture. As a result, a plethora of research investigating its applicability in diverse sectors such as animal husbandry, bioenergy, waste management, pharmaceuticals and food applications have been undertaken. As the body of literature proliferates, a comprehensive examination is essential to reconcile the diverse and sometimes divergent findings across these multiple domains. This section endeavours to synthesize relevant studies that illuminate the multi-dimensional utility of tuber crop skins. This involves critically evaluating both their promising contributions to sustainable development and any existing constraints or limitations. By dissecting the intricacies that define the role of tuber crop skins within the circular agriculture framework, this synthesis aims to offer researchers, agricultural practitioners, and policymakers a detailed perspective for future investigations and sustainable applications.

From waste to wealth: multidisciplinary applications of agricultural peels in circular agriculture

In the domain of circular agriculture, the valorization of yam peels and related biomass appears as a promising avenue for diverse applications, thereby contributing to waste minimization and resource optimization. Initiating the discourse, Aguihe et al. (2022) examined the egg-laying qualities of Japanese quails fed yam peel meal-based diets enriched with enzyme cocktails and yeast. Although their utilization of yam peel was minimal, their study showcases the feasibility of using fermented castor oil seeds to supplement poultry feed. This resultant outcome points toward the broader potential for utilizing alternative bio-resources in animal husbandry, involving yam peels. This idea aligns with circular agriculture's goal of closing the loop by reintegrating waste products back into the production system. A study by Amenaghawon et al. (2023) elevates this concept by employing machine learning models like artificial neural networks (ANN) and adaptive neuro-fuzzy inference systems (ANFIS) to optimize the biotechnological conversion of yam peels for citric acid production. This approach not

only provides a sustainable route for waste valorization but also underscores the role of advanced computational methods in enhancing process efficiencies within a circular agricultural framework. In a related vein, Shao et al. (2022) delve into the extraction and purification of polysaccharides from Chinese yam peel, highlighting their antioxidant and immunomodulatory properties. This suggests that waste products, often disposed, can have high-value applications in functional foods and pharmaceuticals. Onu et al. (2022) extends this circularity paradigm to the energy sector, using thermo-chemical pre-treatment to prepare yam peel substrates for biogas production. Their utilization of machine learning algorithms, like ANN and ANFIS, for process optimization also provides an interdisciplinary approach in actualizing the objective of circular agriculture. The study underscores that the residues from biogas production could be applied as soil conditioners. Thereby bridging the agricultural loop. Li et al. (2021) takes another direction by focusing on the anticorrosion properties of yam peel constituents. This eco-friendly approach towards corrosion prevention has broader implications in reducing the industrial reliance on synthetic chemicals. Thus, offering an avenue for the circular use of agricultural waste for industrial applications.

Again, in the growing field of circular agriculture, the valorization of potato peel waste (PPW) has also emerged as a focal point for researchers who seek to create sustainable and valuable bio-products. With an increasing awareness of the principles of circular economy and the pressing need to mitigate environmental impact, several ground-breaking studies have shed light on how PPW can be upcycled into myriad applications. This encompasses applications ranging from animal feed to green energy, and from waste remediation to functional fillers for food packaging. A study by Almeida et al. (2023) elucidate the paradigmatic shifts required for the valorization of potato peels within the circular economy framework. They offer an in-depth evaluation that sets the stage for subsequent research endeavors. Their work serves as an essential backdrop for understanding the multifaceted potential of PPW, particularly in the extraction of bioactive compounds and as a source of renewable energy. Nematdživa et al. (2022) further expand the horizon by studying the catalytic utilization of PPW for the enzymatic synthesis of bioactive compounds such as phellinsin A. This synthetic approach showcases the utility of PPW as a biotechnological raw material, linking it to health and pharmaceutical applications (another avenue where waste could be repurposed in a circular economy). Concurrently, Saeed et al. (2022) showcases the antioxidant capacity of potato peel extract, underlining its efficacy in stabilizing sunflower oil (a daily consumable prone to oxidative spoilage). This study notably correlates with the principles of circular agriculture by targeting food preservation through waste-derived natural additives. Thus, reducing the necessity for synthetic, often non-biodegradable chemicals. Shifting the lens to green chemical engineering, Daimary et al. (2022) conducted a comprehensive study on the thermochemical conversion of PPW into bio-oil and biochar. The biochar, enriched with alkali and alkaline elements, was subsequently calcined to serve as a heterogeneous catalyst for biodiesel production. This approach stands as a cornerstone in illustrating how circular agriculture could intersect with green chemistry to mitigate waste while generating energy. Importantly, their research offers an innovative yet practical pathway for the transformation of a single agricultural by-product into multiple value-added materials, embodying the very essence of circular agriculture.

Shifting the focus on taro peel starch (TPS) into the central theme of circular agriculture has further manifested the potential of how plant-based waste materials can contribute to sustainability. This is showcased by the study conducted by Bidari et al. (2023). They explored the innovative use of taro peel starch as a renewable material for creating biodegradable films. This pioneering work employs response surface methodology (RSM) to opti-

mize the film's physico-mechanical and water barrier properties. Consequently, their work demonstrates the potential of TPS in replacing petroleum-based components in plastic films. The films exhibited significant attributes such as positive linear effects in thickness, opacity, and water vapor permeability. Moreover, the films showed complete degradation in simulated environmental conditions. Some of which include river water and composting soil, within just five days, signalling its potential as a sustainable alternative for food packaging. When viewed through the lens of circular agriculture, this study complements existing literature by expanding the scope of plant-based waste utility. Also, like potato peel waste, taro peel waste has shown its potential to serve as a bio-based feedstock for sustainable applications. Moreover, the biodegradable films made from TPS underscore the aim of circular agriculture by reducing waste, lowering reliance on non-renewable resources, and contributing to ecological sustainability. As the film degrades quickly, it can be reincorporated into the natural environment, thereby closing the loop in the product life cycle.

The potential of cassava peel has gained notable recognition within the context of circular agriculture, emerging as a prominent sample of tuber skin utilization. For instance, the study by Aladegboye et al. (2022), which explores the use of cassava peels as an asphalt binder, sees a cross-sectional collaboration between civil engineering and agricultural waste management. The study adds a new dimension to the use of agricultural residues, suggesting cassava peels as a sustainable binder in asphalt production. Thus, reducing the reliance on petroleum-based products. Advancing the discourse, the study by Aisien and Aisien. (2020), used cassava peels to produce biogas, serving as an alternative source of renewable energy. In the context of circular agriculture, this suggests a seamless transition from food production to energy generation without incurring waste. Again, the study by Ospino et al. (2022) delve into the utilization of cassava peels for the synthesis of activated carbon. This study capitalizes on the porous structure and high carbon content of cassava peels, re-engineering waste into a cost-effective and efficient material for energy storage devices. In the same vein, Echesi et al. (2022) found that cassava peels can be used as a substrate for pectinase production. Therefore, showcasing its usability in various industrial applications, such as juice extraction and textile processing, thus keeping the agricultural waste in a continuous loop of utility. Contributing to the discourse on bioenergy, Osemwengie et al. (2020) highlighted the use of cassava peels for bioethanol production. Their work presents how this agricultural by-product can be a multifaceted player in the bioenergy sector. Meanwhile, studies by Mourão et al. (2021) and Phetrungnapha et al. (2023) elevate cassava peels from a simple waste material to a complex bioresource capable of yielding polyhydroxybutyrate (PHB) and adsorbents for free fatty acid removal from waste cooking oil, respectively. The former demonstrates the high commercial value of PHB as a biomaterial, while the latter suggests a novel application for cassava peels in the purification of waste cooking oil. Hence, reinforcing the cyclical nature of agricultural waste usage. Continuing this narrative, Kumar et al. (2021) demonstrated the utility of cassava peels as a coagulant aid for institutional wastewater treatment. The techno-economic analysis (TEA) provided by the authors indicates that not only is this approach efficient but also economically viable. Finally, the study by Pulungan et al. (2020) bridges the gap between waste management and material science by optimizing the production of biodegradable plastic from cassava peels. This particular application revolutionizes how we view plastics and their impact on the environment.

In general, the emerging domain of circular agriculture, the diversified valorization of agricultural byproducts such as yam peels, potato peel waste (PPW), taro peel starch (TPS), and cassava peels are attaining academic and industrial attention for their profound implications on sustainability (Oghenejoboh et al. 2021; Khanal et al.

2023; Yafetto et al. 2023). The collective work of studies elucidates a multi-dimensional approach to agricultural waste. This ushers its role from mere dispose to that of a resource capable of complementing energy, feed, healthcare, and material science sectors. Groundbreaking computational methods, chiefly artificial neural networks and adaptive neuro-fuzzy inference systems, are deployed for optimizing biotechnological conversion processes, exemplifying a transdisciplinary mesh between computer science and agriculture. In animal husbandry, waste such as yam and potato peels have shown viability as enriched feeds, offering a sustainable alternative to conventional feedstocks. The potential of these byproducts in producing high-value compounds: antioxidants, polysaccharides, bioactive compounds, and even coagulants for wastewater treatment establishes a precedent for their inclusion in pharmaceuticals and healthcare. Biodegradable films from TPS and plastics from cassava peels point towards an eco-friendly shift in material science. Hence, resonating with circular agriculture's objective of reducing waste and environmental impact. Furthermore, the use of byproducts like PPW and cassava peels for bioenergy production illuminates a pathway to renewable energy solutions. Additionally, another significant aspect of these studies resides in their ability to harmonize waste valorization with contemporary technological advancements. As a result, they foster a sustainable loop that synergizes waste management, resource optimization, and ecological stewardship. The predominant narrative promotes a paradigm shift, encouraging a transition from linear to circular agricultural models. This underscores the imperative to shift our perspective on waste, seeing it not as an endpoint, but rather as the inception of new cycles that offer utility and foster the creation of value. Table 2 presents an overview of the salient findings derived from the relevant studies.

Findings and discussion

In this section, we offer a comprehensive synthesis of pivotal findings derived from rigorously peer-reviewed scholarly contributions focused on the utilization of tuber crop skins in circular agriculture. Each selected manuscript was subjected to a meticulous content analysis that employed advanced statistical methods to unveil emerging trends and discernible patterns within this multifaceted field. Alongside quantitative inquiries, an exhaustive qualitative review was carried out to highlight detailed thematic dimensions embedded in the existing body of work. Expanding upon these diverse lines of analysis, we elucidate the multifunctional utility of tuber crop skins in various sectors such as food application, bioenergy production, animal feed supplementation (animal husbandry), material science and wastewater treatment applications. This analytic framework serves as a lens to unpack the intricate interconnections and contributing factors that underlie the role and valorization of tuber crop skins within the circular agriculture paradigm.

Statistical analysis

In our comprehensive scrutiny of the multifaceted applications of tuber crop skins in circular agriculture, we examined 26 empirical studies, classifying them into five salient thematic clusters: animal husbandry, bioenergy production, material science, wastewater treatment, and food applications. A significant focus was observed in material science, with 10 studies highlighting a keen interest in leveraging tuber crop skins in sustainable material development. This was followed by six studies in the bioenergy sector, reinforcing the growing emphasis on renewable energy sources within the context of circular agriculture.

Table 2. Summary of the relevant literature on the utilization of tuber crop skins in circular agriculture.

Authors Names	Tuber Crop Skin	Area of circular utility	Key Findings
Aisien and Aisien (2020)	Cassava	Bioenergy Production	Cassava peels waste, mixed with cow dung, demonstrated feasibility for biogas and biofertilizer production. The study found that pretreatment with NH_4C^1 led to the best biogas and methane yield. The digester sludge post-digestion exhibited promising biofertilizer qualities, highlighting cassava peels' potential in sustainable energy and fertilizer production.
Candra et al. (2020)	Cassava	Food Applications	The study examined the physicochemical properties of three cassava varieties and their suitability for diet snack chips. Variations in peel thickness, crude fiber, and ash content were significant. Sensory tests showed Pacar cassava peels were preferred for their crispiness and taste, underscoring their potential in diet chip production.
Oladele et al. (2020)	Cassava	Material Science	Chemically treated palm kernel shell fiber (PKSF) and particulate cassava peel (PCP) enhanced epoxy composites' mechanical properties and wear resistance. Specifically, composites with 6% PCP exhibited increased stiffness and deformation resistance, showcasing cassava peel's viability as a sustainable reinforcement in composite production.
Pulungan et al. (2020)	Cassava	Material Science	The study optimized the production of biodegradable plastic from cassava peel starch. It identified optimal conditions for drying temperature and duration, resulting in a bioplastic with good mechanical properties and biodegradability. The research highlights cassava peel's potential as a sustainable source for bioplastic production, offering an eco-friendly solution to plastic waste.
Azizi et al. (2021)	Potato	Food Applications	The study showcased the development of an optimized potato snack using potato skins, achieving dual objectives: minimizing food waste and providing a healthier snack option. This innovation highlights the potential of potato skins in producing sustainable, nutritious food products, emphasizing their value in eco-friendly and health-oriented food solutions.
Castro et al. (2021)	Cassava	Material Science	The study produced polyhydroxyalkanoates (PHAs) from cassava peel waste using <i>Cupriavidus necator</i> , identifying optimal production conditions. The biopolymer was then transformed into microfibers via electrospinning, showcasing cassava peel waste's innovative application in biopolymer manufacturing, with potential industrial and biomedical uses.
Kumar et al. (2021)	Cassava	Wastewater Treatment	Cassava peel starch (CPS) was evaluated as an eco-friendly coagulant for institutional wastewater treatment. The study optimized the treatment process using CPS and found it to be economically viable with significant removal efficiencies for turbidity, TSS, and COD. CPS proved to be a sustainable alternative to chemical coagulants, with potential to contribute to wastewater management and pollution reduction.
Li et al. (2021)	Yam	Material Science	Chinese yam peel extracts (CYPE), containing diosgenin, batatasin-I, batatasin-III, and yam polysaccharide, effectively prevent copper corrosion in artificial seawater. Acting as mixed-type inhibitors with mainly anodic effects, CYPE's high efficiency at elevated temperatures positions it as a potential eco-friendly solution for protecting copper in marine settings.
Mourão et al. (2021)	Cassava	Material Science	Using cassava peel hydrolysate, <i>Stigeoclonium</i> sp. B23 produced polyhydroxybutyrate (PHB), a promising biomaterial, achieving 12.16% PHB content. Despite some toxicity in zebrafish embryos from PHB nanoparticles, the study highlights cassava peel as a renewable source for PHB production, presenting an alternative for biomaterial applications and tackling agro-industrial waste.
Papathoti et al. (2021)	Cassava	Bioenergy Production	The study focused on boosting ethanol production from cassava peel waste through alkali-assisted hydrothermal pretreatment. Using the thermotolerant <i>Kluyveromyces marxianus</i> , higher ethanol yields (0.44 g/g) and efficiency (86.11%) were achieved with simultaneous saccharification and fermentation (SSF) over separate hydrolysis and fermentation (SHF), highlighting cassava peel's viability as a bioethanol feedstock.

Accepted manuscript (author version)

Aguihe et al. (2022)	Yam	Animal Husbandry	Feeding Japanese quails with diets including yam peel meal, enzymes, and yeast showed varied egg quality effects. Adding yam peel, FCSM, and methionine types improved health and productivity. Yet, high yam peel and FCSM levels (up to 150 g/kg) reduced carcass weight without harming other physiological traits. This indicates yam peel's potential in boosting egg quality, despite carcass weight considerations.
Aladegboye et al. (2022)	Cassava	Material Science	The study assessed Cassava Peel Ash (CPA) as an asphalt mixture modifier, focusing on sustainable pavement material development and agricultural waste reduction. Findings revealed that CPA concentrations notably affected the mixtures' volumetric properties, suggesting CPA's efficacy in improving asphalt performance.
Daimary et al. (2022)	Potato	Bioenergy Production	Potato peels were transformed into biochar, bio-oil, and a biodiesel catalyst, with peak yields of 23.60% bio-oil and 29.50% biochar at 500°C. The potassium-rich catalyst achieved a 97.50% oil conversion in biodiesel production, showcasing potato peels' sustainable conversion into valuable fuel and chemical products.
Ebrahimian et al. (2022)	Potato	Bioenergy Production	The study outlined the valorization of potato peel waste (PPW) for producing biofuel, bioplastics, and biosorbents, showcasing its potential in sustainable bioproduct development via biological and thermochemical methods. This emphasizes PPW's importance in a PPW-based biorefinery approach.
Haque et al. (2022)	Potato	Food Applications	Ethanol-extracted antioxidants from potato peels showed 89.14% radical scavenging activity, presenting a natural solution for food preservation. Incorporating a 3% ethanol extract into nacho dough enhanced antioxidant activity, indicating a sustainable method to boost food product nutrition using potato peels.
Min et al. (2022)	Potato	Material Science	Potato skin carbon dots, used as nanofillers in gelatin-based packaging, enhanced the material's antioxidant and antimicrobial activities, water vapor permeability, hydrophobicity, UV-barrier properties, and mechanical stability. This underscores their potential as sustainable, value-added nanofillers for packaging uses.
Nemadziva et al. (2022)	Potato	Material Science	Potato peels were utilized to generate chlorogenic acid (CLA) and caffeic acid (CFA), which laccase converted into phellinsin A, boosting antioxidant capacity with low cytotoxicity. This indicates their potential as antioxidant agents across applications, highlighting potato peels' effectiveness in producing valuable bioactive compounds via laccase catalysis.
Onu et al. (2022)	Yam	Bioenergy Production	Optimizing yam peel thermo-chemical pretreatment for biogas through ANFIS, ANN, and RSM models showed ANFIS's superior accuracy. Optimal conditions led to a biogas yield with 92% methane. The research underscores yam peels' viability for biogas production and suggests using digestion sediments in agriculture.
Ospino et al. (2022)	Cassava	Material Science	The study involved producing activated carbon from cassava peels for use as electrode material in supercapacitors. The activated carbon, treated with potassium hydroxide and phosphoric acid, showed a specific capacitance of 64.18 F/g, indicating its viability as a cost-effective and sustainable electrode material for energy storage applications.
Saeed et al. (2022)	Potato	Food Applications	Potato peel extract, prepared with hexane, exhibited strong antioxidant activity, on par with synthetic antioxidants like BHA, in stabilizing sunflower oil. Sunflower oil enhanced with this extract showed increased oxidative stability, underscoring potato peel extract's potential as a natural antioxidant for extending vegetable oils' shelf life.

Accepted manuscript (author version)

Singh et al. (2022)	Potato	Wastewater Treatment	Potato peel biochar outperformed potato peel powder in adsorbing chlorpyrifos, achieving a 72.06% removal efficiency. This highlights the potential of potato peel waste as an economical and sustainable water treatment adsorbent.
Shao et al. (2022)	Yam	Food Applications	Optimized extraction of Chinese yam peel polysaccharide (CYPP) mainly yielded glucose and galactose. CYPP-1 showed antioxidant, immunomodulatory, and hypoglycemic effects, indicating its potential for functional food and pharmaceutical industries.
Amenaghawon et al. (2023)	Yam	Bioenergy Production	Using yam peels with EDTA (0.3 g/l) and coconut oil (5.0% w/w) significantly boosted citric acid production by 49.1%, achieving 43.08 g/l. This showcases the yam peels' biochemical potential when combined with specific stimulants, although sodium fluoride lacked a stimulatory effect. The study reveals yam peels' promising role in citric acid bioproduction.
Bidari et al. (2023)	Taro	Material Science	Taro peel starch (TPS) was used to create biodegradable films via casting, with optimized TPS and glycerol levels impacting film characteristics. These TPS films, exhibiting an amorphous-type crystallinity, fully degraded in 5 days under environmental conditions, confirming their viability as sustainable food packaging polymers.
Goyi et al. (2023)	Potato	Wastewater Treatment	Potato peel hydrochar, created via hydrothermal carbonization, demonstrated enhanced Congo red adsorption over raw potato peel powder. This highlights its viability as a sustainable water treatment adsorbent, aligning with pseudo-second-order kinetics and Freundlich isotherm models.
Phetrungnapha et al. (2023)	Cassava	Wastewater Treatment	Cassava peel-derived adsorbent effectively removed free fatty acids from waste cooking oil. The optimal adsorbent, developed through calcination and NaOH modification, demonstrated high adsorption capacity and was characterized using FT-IR and SEM. The study revealed that this adsorbent is a promising, sustainable solution for purifying waste cooking oil, highlighting cassava peels' utility in environmental remediation.

Wastewater treatment and food applications were explored in four and five studies respectively, pointing to an evolving interest in these domains for environmental management and culinary innovation. Notably, a single study addressed animal husbandry, suggesting a nascent yet promising area for further exploration. These categories outline the current scope of research and underscore areas for future scholarly exploration. They offer insights into untapped potential and encourage a multidisciplinary approach to circular agriculture. Fig. 2 graphically encapsulates these five pivotal thematic clusters.




Figure 2. Utilization of tuber crop skin in circular agriculture.

Again, to further examine the evolving trend in the utilization of tuber crop skins within circular agriculture over time, we conducted a chronological analysis of academic publications. The time frame of this scrutiny extended from 2020 to 2023. During this period, there was a notable rise in the number of publications: starting with four articles in 2020, rising to six in 2021, and then further increasing to twelve studies in 2022, before experiencing a decrease to four by the third quarter of 2023. This dynamic pattern may be indicative of several underpinning factors. The modest beginnings in 2020 possibly reflect the growing stage of this research avenue. The spike in 2022 could signify an academic recognition of the urgency for sustainable agriculture, particularly in the wake of increasing environmental concerns. Conversely, the dip in 2023, despite being only partial data for the year, may point towards a shift in research concentrations or possibly a period of academic digestion and integration of existing studies. This temporal examination deepens our understanding of the evolving landscape of this field. Furthermore, it highlights the growing importance of utilizing tuber crop skins within the broader contexts of circular agriculture and sustainability.

Additionally, our study aimed to clarify the geographical spread of scholarly research on the utilization of tuber crop skins in circular agriculture. Our analysis identified empirical works from 14 nations, with a predominant focus on Africa, South America, and Asia. Intriguingly, Asia accounted for a significant portion of the research collection. This trend may be linked to the region's growing dedication to agricultural sustainability and technological innovation. Africa followed closely in significance, with a notable presence in the research. This may be indicative of the continent's increasing emphasis on sustainable agricultural practices as a means to address persistent challenges such as food insecurity and climate change adaptation. In contrast, South America's involvement lagged behind both Asia and Africa. This could signify either an early stage of scholarly inquiry in the region or a comparatively lesser emphasis on tuber crop skins in the broader scope of agricultural research. Remarkably, an evident absence of research contributions was noted from Europe, Australia and North America. This void is particularly striking given these regions' advancement in agricultural technologies and sustainability paradigms. The dearth of academic output from these continents suggests a couple of possibilities. Firstly, it could indicate that the theme of tuber crop skin utilization in circular agriculture has not yet gained substantial traction in academic discourse within these regions. Secondly, it may reflect that other priorities in agricultural sustainability are currently taking precedence over this specific area of focus. This geographical gap in academic contributions adds complexity to our understanding, highlighting areas of deficiency that present opportu-

nities for future scholarly research. This geospatial analysis provides a comprehensive overview of current academic trends and raises important questions about the disparity in scholarly activity across different continents. Fig. 3 captures this geographical diversity, thereby enriching our overall understanding of the global research landscape in this specialized domain of circular agriculture.




Figure 3. Regions of empirical investigation on the utilization of tuber crop skins in circular agriculture.

Multifaceted applications of tuber crop skins in circular agriculture

The use of tuber crop skins in circular agriculture has garnered considerable attention from both the academic and industrial sectors. This has led to a multidisciplinary dialogue involving agronomists, environmental scientists, and policymakers. In this section, we present a comprehensive synthesis of existing research, examining the versatile applications of tuber crop skins in the field of circular agriculture. Based on a detailed analysis of the relevant peer-reviewed articles, the study affirms the considerable potential of tuber crop skins as sustainable resources for a variety of sectors. These skins find application across various sectors, including animal husbandry, bioenergy production, material science, wastewater treatment, and food applications. These skins have proven to be invaluable, serving as nutrient-rich feed in animal husbandry and as promising feedstocks for bioenergy production. Their high content of cellulose, hemicellulose, and lignin offers groundbreaking opportunities in material science. Additionally, their absorptive properties present viable solutions for wastewater treatment. Moreover, their nutrient-dense profiles are well-suited for food fortification and the creation of functional foods. This comprehensive assessment seeks to present a detailed overview of the scope and distribution of tuber skin applications across the identified sectors, as presented in Table 3.

Table 3. Content analysis of selected articles showcasing the utilization of tuber crop skins in circular agriculture across varied domains.

Authors Names	Animal Hus- bandry	Bioenergy Pro- duction	Material Science	Wastewater Treatment	Food Ap- plications
Aisien and Aisien (2020)		✓			
Candra et al. (2020)					✓
Oladele et al. (2020)			✓		
Pulungan et al. (2020)			✓		
Azizi et al. (2021)					✓
Castro et al. (2021)			✓		
Kumar et al. (2021)				✓	
Li et al. (2021)			✓		
Mourão et al. (2021)			✓		
Papathoti et al. (2021)		✓			
Aguihe et al. (2022)	✓				
Aladegboye et al. (2022)			✓		
Daimary et al. (2022)		✓			
Ebrahimian et al. (2022)		✓			
Haque et al. (2022)					✓
Min et al. (2022)			✓		
Nemadziva et al. (2022)			✓		
Onu et al. (2022)		✓			
Ospino et al. (2022)			✓		
Saeed et al. (2022)					✓
Singh et al. (2022)				✓	
Shao et al. (2022)					✓
Amenaghawon et al. (2023)		✓			
Bidari et al. (2023)			✓		
Goyi et al. (2023)				✓	
Phetrungnapha et al. (2023)				✓	

Implication and recommendation of the multifaceted applications of tuber crop skin in circular agriculture

The multifaceted applications of tuber crop skins within circular agriculture systems present a promising avenue for sustainable resource management, energy production, material innovation, and food security. This section outlines the implications and recommendations stemming from the utilization of these agricultural by-products across various domains. From enhancing animal husbandry practices and advancing bioenergy solutions to pioneering in material science innovations and unlocking nutritional potential in food applications, tuber crop skins are an untapped resource with considerable environmental and economic benefits. This section highlights and discusses issues, solutions, and implications regarding the use of tuber crop skins, while also suggesting areas for further exploration.

Sustainable animal husbandry

Exploring the use of tuber crop skins in animal husbandry presents a promising solution within the framework of circular agriculture. These skins, rich in essential nutrients like carbohydrates, fibers, and minerals, offer a sustainable alternative to conventional animal feedstocks such as corn and soy, potentially reducing reliance on resource-intensive feeds (Rahman et al. 2021; Okike et al. 2022; Rahmani et al. 2022). The inherent organic matter in tuber crop skins enhances livestock gut health and well-being, contributing to improved quality of animal-derived products (Wanapat et al. 2013; Ruqayyah et al. 2014). Furthermore, their fibrous nature provides environmentally friendly bedding materials, showcasing a multifaceted approach to sustainability (Omole et al. 2013; Gebrechristos and Chen 2018). The potential health benefits from bioactive compounds in these skins, such as antioxidants, further posit an opportunity to enhance animal health and productivity (Surai 2014; Oloruntola 2020). This integration not only supports waste reduction and resource optimization but also aligns with the broader goals of circular agriculture by closing nutrient loops and enhancing the ecological footprint of livestock farming. Additionally, it offers socio-economic benefits, particularly for smallholder farmers, by providing a cost-effective alternative to traditional feeds and bedding materials, thus enhancing economic resilience (Kalio et al. 2015). Despite the evident potential, there is a notable gap in empirical research focusing on this application, highlighting the need for comprehensive studies to assess the nutritional, health, environmental, and economic impacts of incorporating tuber crop skins into animal husbandry. This area represents a promising avenue for future research, with the potential to redefine conventional practices in animal husbandry by promoting more sustainable, integrated approaches within circular agricultural systems.

Bioenergy innovation in with tuber crop skins

Within the circular agriculture frameworks, the conversion of tuber crop skins to bioenergy stands out as a key area of innovation. Tuber crop skins are increasingly recognized not just as agricultural waste but as valuable bioenergy feedstock for producing biogas, bioethanol, and biodiesel (Nair et al. 2022; Almuhayawi et al. 2023). The rich fibrous and starchy nature of tuber crop skins offers an abundant source of fermentable sugars (Okekunle et al. 2016; Awogbemi et al. 2022). This makes them ideal for bioenergy production processes. Anaerobic digestion yields methane-rich biogas, while fermentation processes produce bioethanol (Wu et al. 2016). Advances in pyrolysis and gasification have further expanded their use, facilitating the creation of syngas and bio-oil (Foong et al. 2023). The transformation of tuber crop skins into bioenergy aligns with sustainable energy goals, offering a renewable alternative to diminish reliance on fossil fuels. This shift not only addresses the pressing need for sustainable energy solutions but also integrates waste management into energy production, enhancing agricultural circularity (Awogbemi et al. 2022). The holistic use of tuber crop skins in bioenergy production exemplifies the circular economy's ethos, turning potential waste into valuable energy resources and contributing to environmental sustainability (Wu et al. 2016). The need for cross-disciplinary research and policy support is highlighted by this multifaceted approach. It is essential to unlock the full potential of tuber crop skins in bioenergy within circular agriculture. This promises a path towards energy security and environmental sustainability.

Material science innovations with tuber crop skins

The integration of tuber crop skins into material science represents a significant advancement within circular agriculture, offering a sustainable solution to the challenges of waste management and environmental degradation (Borah et al. 2017). Recognized for their rich cellulose, hemicellulose, and lignin content, tuber crop skins are becoming essential

raw materials for developing bio-composites, bio-based plastics, and functional films (Bello et al. 2023). These applications not only enhance the mechanical properties and biodegradability of materials but also provide eco-friendly alternatives to synthetic plastics, aligning with efforts to reduce environmental pollution (Sugumaran et al. 2015; Versino et al. 2015; Charles et al. 2022). Further exploring their utility, the bioactive compounds in tuber crop skins are being harnessed for antimicrobial and antioxidant properties, making them ideal for food packaging that demands extended shelf life and microbial resistance (Miller et al. 2022; Thuppahige et al. 2023). Their application in aerogels and adsorbents for environmental remediation, targeting pollutants and heavy metals in wastewater, and even in energy storage devices like supercapacitors, showcases the breadth of their potential impacts (Osman et al. 2019; Nathan et al. 2021; Minisha et al. 2022; Odeyemi et al. 2023). This transformation of tuber crop skins from waste to valuable material science assets exemplifies the principles of the circular economy, emphasizing the creation of sustainable materials from agricultural by-products. The convergence of waste reduction, innovative material production, and environmental stewardship underlines the holistic benefits of leveraging tuber crop skins in circular agriculture. This shift necessitates focused research and supportive policy frameworks to fully exploit their sustainability potential, marking a pivotal step towards realizing circular agricultural systems.

Wastewater treatment innovations with tuber crop skins

The role of tuber crop skins in wastewater treatment within circular agriculture systems presents an innovative approach to sustainable resource management (Bhatnagar et al. 2015; Kumar et al. 2017). These skins, abundant in organic matter such as carbohydrates and proteins, exhibit natural adsorptive properties for removing pollutants, offering a low-cost alternative to traditional adsorbents (Kumar et al. 2017; Asharuddin et al. 2018; Nahas et al. 2019; Rodríguez et al. 2021). Processed into biochar or activated carbon, tuber crop skins effectively eliminate a variety of contaminants, matching the efficiency of commercial adsorbents due to their high surface area and functional groups (Asuquo et al. 2018; Gale et al. 2021; Geça et al. 2022; Liang et al. 2015). Moreover, the enzymatic potential of tuber skins in biodegradation processes highlights their ability to act as biological catalysts, broadening the spectrum of treatable pollutants, including pharmaceuticals and synthetic dyes (Belcaid et al. 2023; Hoseinzadeh et al. 2014; Kayiwa et al. 2022). This dual functionality aligns with waste reduction goals and enhances water recycling in agriculture. It reinforces circular economy principles by enabling the reuse of treated wastewater for irrigation (Abdelraof et al. 2019). The underexplored potential of tuber crop skins in wastewater treatment emphasizes the necessity for in-depth research. This research should focus on standardizing treatment methods and maximizing efficiency. The goal is to integrate tuber crop skins more effectively into sustainable agriculture and environmental conservation efforts. Emphasizing their multifaceted utility in wastewater treatment reinforces circular agriculture's commitment to environmental and economic sustainability.

Nutritional innovations with tuber crop skins

In the aspect of circular agriculture, tuber crop skins offer more than just a solution for waste management; they play a pivotal role in food innovation, enhancing nutritional content and facilitating the development of functional foods (Samotyja 2019; Vaitkevičienė 2019; Triwidatin and Djuanda 2023). These often-overlooked by-products, recognized for their rich composition of vitamins, minerals, fiber, and bioactive compounds like phenols and flavonoids, are valuable for enriching a variety of food products. From bakery goods to beverages and snacks, they showcase a sustainable approach to food production (Curti et al. 2016; Ben et al. 2017). The functional attributes of tuber skins, such as their antioxidant and colorant properties, provide natural alternatives to synthetic additives, enhancing food safety and nutri-

tional profiles (Javed et al. 2019; Helal et al. 2020). Moreover, their starch and fiber content offers potential as natural thickeners or stabilizers in culinary applications, contributing to innovative food textures and consistencies (Rengsutthi and Charoenrein 2011; Xie et al. 2018). Addressing global challenges like food security and malnutrition, the strategic use of tuber crop skins in food products underscores their societal and nutritional significance, especially in regions facing food scarcity (Alemu 2020). The transformation of tuber skins into high-value products such as protein concentrates and dietary fibres on an industrial scale aligns with the goals of sustainable food systems (Caliskan et al. 2023). This process provides eco-friendly and economically viable alternatives (Singh et al. 2023). This comprehensive approach not only underscores the necessity of waste reduction and nutrient recycling but also positions tuber crop skins as a cornerstone of sustainable food production. The extensive benefits of tuber skins in food applications underscore the importance of continuous interdisciplinary research. This research should focus on standardizing extraction methods and quantifying bioactive compounds. The goal is to meet regulatory standards and consumer expectations, fully leveraging the potential of tuber crop skins in circular agriculture.

Summary and future research directions in the multifaceted utilization of tuber crop skins in circular agriculture systems

The diverse applications of tuber crop skins substantially enhance the landscape of circular agriculture. They impact crucial domains including animal husbandry, bioenergy production, material science, wastewater treatment, and food applications (Okike et al. 2015; Nair et al. 2022). The potent nutrient profiles, fibrous properties, and bioactive constituents of tuber crop skins enable their use in sustainable and innovative solutions (Hoseinzadeh et al. 2014; Charles et al. 2022). Tuber crop skins serve various roles, from nutritional supplements in animal feed to feedstocks for bioenergy, building blocks in bio-based materials, and agents in wastewater purification and food fortification. In each of these applications, they embody the essence of resource optimization and waste minimization. These are fundamental goals of circular agriculture. Despite the apparent robustness of these applications, the use of tuber crop skins in circular agriculture is still relatively underexplored and fragmented. A notable deficiency is the lack of interdisciplinary research that melds nutritional science, environmental sustainability, and economic modeling. Such integrated research is crucial for a comprehensive understanding of the impact and feasibility of these applications, as noted in studies by Kumar et al. (2021) and Daimary et al. (2022). Empirical studies focusing on the long-term outcomes, scalability, and the Technological Readiness Level (TRL) of these utilizations are notably lacking. Moreover, research has not yet completely delved into the mechanisms underlying the bioactive properties of tuber crop skins. This includes understanding their roles in enhancing animal health, improving bioenergy conversion efficiency, and aiding pollutant sorption in wastewater treatment. In the area of food applications, quantifiable metrics for bioactive compounds, and regulatory compliance are largely undeveloped. Such limitations underscore the urgent need for robust, methodologically sound, and comprehensive studies to elucidate the complexities involved. Additionally, incorporating tuber crop skins into existing agricultural and industrial systems brings up issues related to supply chain logistics, quality control, and policy frameworks. These critical aspects are yet to be thoroughly addressed in the academic literature. Ethical considerations regarding resource allocation, especially in food-scarce regions, also warrant scholarly attention. Finally, the use of tuber crop skins offers a promising route to enhance the sustainability and functionality of circular agricultural systems. However, the field is at a critical juncture that calls for rigorous empirical research and collaborations among multiple stakeholders. It is through such focused endeavours that the full spectrum of opportunities afforded by tuber crop skins can be unlocked, actualized, and optimized for the future.

Limitations of the study

In reflecting upon the limitations of our study, it is important to acknowledge certain constraints that have influenced the scope and depth of our findings. Our study was primarily confined to publications from 2020 to 2023. While this timeframe allowed for a focused review, it may have excluded both earlier foundational studies and very recent research. This potentially overlooks broader historical insights or cutting-edge developments in the field. Additionally, the geographical representation in the reviewed literature was uneven, potentially introducing a regional bias. This disparity might affect the interpretation of global trends in the utilization of tuber crop skins. Particularly, certain regions, especially those leading in research and application, could disproportionately influence the study's conclusions. Another limitation is our focus on studies published in English. This language restriction might have inadvertently excluded significant research in other languages. Consequently, narrowing the global perspective and understanding of innovations in circular agriculture involving tuber crop skins. There is also the potential for selection bias. Despite efforts for a comprehensive review, our selection criteria might have favored certain research methodologies or applications. This could result in overlooking divergent or contradictory views, leading to a somewhat restricted interpretation of the current state of the field. Moreover, our reliance on academic databases (ScienceDirect and WoS) for sourcing articles means that our review is subject to publication biases and the predominance of certain journals. This could skew the representation of specific aspects of research in the utilization of tuber crop skins. Lastly, our focus predominantly on academic literature may not fully capture the practical advancements and challenges in the industry. Some innovative applications or real-world implementations of tuber crop skins in circular agriculture might not be extensively documented in scholarly articles. Recognizing these limitations is essential for properly contextualizing the study's findings. These constraints should be considered when applying our insights to future research, policy-making, or practical applications in the aspect of circular agriculture and sustainable use of agricultural by-products.

Conclusion

This study highlights the transformative role of tuber crop skins in circular agriculture, marking a shift toward sustainable food production and resource management. The study has explored their multifaceted utility, from serving as nutrient-rich feed supplements in animal husbandry to emerging as valuable feedstocks for bioenergy production. In material science, their substantial cellulose, hemicellulose, and lignin content make them ideal for bio-composites and bio-based plastics. This aligns with the waste-to-resource ideology at the heart of circular agriculture. Their potential is further demonstrated in wastewater treatment and food applications, where they help treat pollutants and enrich food products, addressing global challenges such as malnutrition. However, despite these promising areas, there is a lack of comprehensive empirical research on the long-term environmental, technical, and economic implications of these applications. The study highlights this gap as a crucial area for future interdisciplinary research. Such research is necessary to fully realize the potential of tuber crop skins in a sustainable, economically viable, and resource-efficient agricultural paradigm. This study sets the stage for future explorations, advocating for an integrated approach to address these gaps. In doing so, it underscores the importance of tuber crop skins in advancing global sustainability goals and contributing to a restorative agricultural process.

Funding: This research was supported by the Integrated Research Management Centre (IRMI), Universiti Teknologi MARA (UiTM), under Grant number 600-RMC/GIS 5/3 (005/2023).

Author contributions: Aziz Caliskan: Conceptualization, Methodology, Visualization, Writing-Original draft preparation and Revising of manuscript; Norhidayah Abdullah: Supervision, Funding acquisition, Editing, Visualization and Proof-reading of manuscript; Noriza Ishak: Supervision, Methodology, Visualization, Validation and Proof-reading of manuscript; Divine Senanu Ametefe: Writing-Original draft preparation, Methodology, validation, Editing and Proof-reading of manuscript, Visualization, Validation and Revising of manuscript; Imene Tatar Caliskan: Curation of Literature, Methodology, validation, Visualization, Editing and Revising of manuscript.

References

- Abdelraof M, Hasanin M, El-Saied H (2019) Ecofriendly green conversion of potato peel wastes to high productivity bacterial cellulose. *J Carbpol* 211: 75–83. <https://doi.org/10.1016/j.carbpol.2019.01.095>
- Abioye S, Oyedele L, Akanbi L, Ajayi A, Davila J, Bilal M, Akinade O, Ahmed A (2021) Artificial intelligence in the construction industry: A review of present status, opportunities and future challenges. *J Jobe* 44: 315–329. <https://doi.org/10.1016/j.jobe.2021.103299>
- Aguihe P, Kehinde A, Joshua D, Ilaboya I, Samuel K, Odu O (2022) 34. Egg laying qualities of Japanese quails fed yam peel meal based diets with enzyme cocktail+yeast supplementation. *J Anscip* 13(1): 23–24. <https://doi.org/10.1016/j.anscip.2022.03.035>
- Aisien F, Aisien E (2020) Biogas from cassava peels waste. *Detritus* 10(6): 100–108. <https://doi.org/10.31025/2611-4135/2020.13910>
- Aladegboye O, Oguntayo O, Al-Ihekwa E, Daniel T, Chiadighikaobi P, Ng P (2022) Evaluation of volumetric properties of cassava peel ash modified asphalt mixtures. *Eng J* 8(10): 20120–20124. <https://doi.org/10.28991/CEJ-2022-08-10-07>
- Alemu E (2020) Malnutrition and its implications on food security. *Zero Hunger* 1(1): 509–518. https://doi.org/10.1007/978-3-319-95675-6_32
- Almeida P, Gando-Ferreira L, Quina M (2023) Biorefinery perspective for industrial potato peel management: Technology readiness level and economic assessment. *J Environ Chem Eng* 11(3): 172–186. <https://doi.org/10.1016/j.jece.2023.110049>
- Almuhayawi M, Hassan E, Alkuwaity K, Abujamel T, Mokhtar J, Niyazi H, Almasaudi S, Alamri T, Najjar A, Zaber-mawi N, Azhar E, Makki R, Niyazi H, Harakeh S (2023) Enzymatic-based hydrolysis of digested potato peel wastes by amylase producing fungi to improve biogas generation. *Catalysts* 13(5): 913–920. <https://doi.org/10.3390/catal13050913>
- Amenaghawon A, Omoruyi B, Kenneth I, Okedi M, Esenogho G, Oyefolu P, Muojama O, Otuya I, Eshiemogie S, Okoh R, Anyalewechi C (2023) Biotechnological conversion of yam peels for enhanced citric acid production: Data-driven machine learning modeling and global sensitivity analysis of the impact of metabolic stimulants. *Ind Crops Prod* 191: 228–41. <https://doi.org/10.1016/j.indcrop.2022.116022>
- Asharuddin S, Othman N, Shaylinda N, Zin M, Tajarudin H (2017) A chemical and morphological study of cassava peel: A potential waste as coagulant aid. *ISCEE* 13(1): 35–49. <https://doi.org/10.1051/mateconf/201710306012>
- Asharuddin N, Shaylinda N, Zin M, Azan Tajarudin H, Fadhil M, Din M, Kumar V (2018) Performance assessment of cassava peel starch and alum as dual coagulant for turbidity removal in dam water. *IJIE* 10(4): 185–192. <https://doi.org/10.30880/ijie.2018.10.04.029>

- Asuquo E, Martin A, Nzerem P (2018) Evaluation of Cd (ii) ion removal from aqueous solution by a low-cost adsorbent prepared from white yam (*dioscorea rotundata*) waste using batch sorption. *Chem Eng* 2(3): 1-35. <https://doi.org/10.3390/chemengineering2030035>
- Awogbemi O, Kallon D, Owoputi A (2022) Biofuel generation from potato peel waste: Current state and prospects. *Recycling* 7(2): 74–90. <https://doi.org/10.3390/recycling7020023>
- Azizi A, Sethi S, Joshi A, Arora B (2021) Utilisation of potato peel in fabricated potato snack. *Potato Res* 64(4): 587–599. <https://doi.org/10.1007/S11540-021-09492-2>
- Belcaid A, Beakou B, Bouhsina S, Anouar A (2023) New insights on manganese dioxide nanoparticles loaded on cellulose-based biochar of cassava peel for the adsorption of three cationic dyes from wastewater. *Int J Biol Macromol* 241: 493–502. <https://doi.org/10.1016/j.ijbiomac.2023.124534>
- Bello T, Eze E, Usman M, Isa M (2023) Characterization of bioplastics produced from yam and potato peels using hydrochloric and acetic acids. *Biomass Convers Biorefin* 1(3): 1–12. <https://doi.org/10.1007/s13399-023-04021-2/figures/12>
- Ben K, Bouaziz F, Zouari-Ellouzi S, Chaari F, Ellouz-Chaabouni S, Ellouz-Ghorbel R, Nouri-Ellouz O (2017) Improvement of texture and sensory properties of cakes by addition of potato peel powder with high level of dietary fiber and protein. *Food Chem* 217: 668–677. <https://doi.org/10.1016/j.foodchem.2016.08.081>
- Bhatnagar A, Sillanpää M, Witek-Krowiak A (2015) Agricultural waste peels as versatile biomass for water purification – A review. *J Chem Eng* 270: 244–271. <https://doi.org/10.1016/j.cej.2015.01.135>
- Bidari R, Abdillah A, Ponce R, Charles A (2023) Characterization of biodegradable films made from taro peel (*Colocasia esculenta*). *Starch Plym* 15(2): 338–355. <https://doi.org/10.3390/polym15020338>
- Borah P, Das P, Badwaik L (2017) Ultrasound treated potato peel and sweet lime pomace based biopolymer film development. *Ultrason Sonochem* 36: 11–19. <https://doi.org/10.1016/j.ultsonch.2016.11.010>
- Cainelli G, D'Amato A, Mazzanti M (2020) Resource efficient eco-innovations for a circular economy: Evidence from EU firms. *Res Policy* 49(1): 1–8. <https://doi.org/10.1016/j.respol.2019.103827>
- Caliskan A, Abdullah N, Ishak N, Tatar, I (2023) Physicochemical, microbial and sensory properties of wild carob bar : A shelf-life study. *Int J Gastron Food Sci* 31(1): 1–7. <https://doi.org/10.1016/j.ijgfs.2023.100668>
- Candra K, Ainudin A, Arifin M, Yuliani Y (2020) Physicochemical characteristics of cassava peel from samarinda and its acceptability for diet snack chips. *Agri Tech* 40(4): 299–305. <https://doi.org/10.22146/agritech.35291>
- Castro O, León E, Arias M, Cesario M, Ferreira F, Fonseca M, Segura A, Valencia P, Simpson R, Nuñez H, Contreras-Calderon J (2021) Characterization and production of a polyhydroxyalkanoate from cassava peel waste: Manufacture of biopolymer microfibers by electrospinning. *J Polym Environ* 29(1): 187–200. <https://doi.org/10.1007/S10924-020-01861-1>
- Charles A, Motsa N, Abdillah, A (2022) A comprehensive characterization of biodegradable edible films based on potato peel starch plasticized with glycerol. *Polym* 14(17): 3462–3476. <https://doi.org/10.3390/polym14173462/s1>
- Curti E, Carini E, Diantom A, Vittadini E (2016) The use of potato fibre to improve bread physico-chemical properties during storage. *Food Chem* 195: 64–70. <https://doi.org/10.1016/j.foodchem.2015.03.092>
- Dahal K, Li X, Tai H, Creelman A, Bizimungu B (2019) Improving potato stress tolerance and tuber yield under a climate change scenario – a current overview. In *Front Plant Sci* 10: 563–579. <https://doi.org/10.3389/fpls.2019.00563>

- Daimary N, Eldiehy K, Boruah P, Deka D, Bora U, Kakati B (2022) Potato peels as a sustainable source for biochar, bio-oil and a green heterogeneous catalyst for biodiesel production. *J Environ Chem Eng* 10(1): 107108–107121 <https://doi.org/10.1016/j.jece.2021.107108>
- Ebrahimian F, Denayer J, Karimi K (2022) Potato peel waste biorefinery for the sustainable production of biofuels, bioplastics, and biosorbents. *Bioresour Technol* 360: 644–658. <https://doi.org/10.1016/j.biortech.2022.127609>
- Echesi S, Sopuruchukwu F, Odu N (2022) Optimization of pectinase production from bacillus subtilis pse-8 using cassava peels as substrate in submerged fermentation through response surface methodology (RSM). *J Adv Microbiol* 22(11): 56–66. <https://doi.org/10.9734/JAMB/2022/v22i11682>
- Elhassaneen Y, Ragab S, Mashal R (2016) Improvement of bioactive compounds content and antioxidant properties in crackers with the incorporation of prickly pear and potato peels powder. *Int J Food Sci Nutr* 5(1): 53–61. <https://doi.org/10.11648/j.ijnfs.20160501.18>
- El-Sawi S, Ibrahim M, Bassuiny R, Merghany R (2023) Antioxidant, cytotoxic and antimicrobial efficacy of potato peels, taro peels, and husk and silk of corn. *Proc Natl Acad Sci* 93(3): 619–626. <https://doi.org/10.1007/s40011-023-01473-4>
- Fernandez J, Gu X, Chen S (2020) Techno-economic assessment of bioactive compound recovery from potato peels with sequential hydrothermal extraction. *J Clean Prod* 282: 124–164. <https://doi.org/10.1016/j.jclepro.2020.124356>
- Foong S, Chan Y, Lock S, Chin B, Yiin C, Cheah K, Loy A, Yek P, Chong W, Lam S (2023) Microwave processing of oil palm wastes for bioenergy production and circular economy: Recent advancements, challenges, and future prospects. *Bioresour Technol* 369: 128–137. <https://doi.org/10.1016/j.biortech.2022.128478>
- Gale M, Nguyen T, Moreno M, Abdulaziz K (2021) Physiochemical properties of biochar and activated carbon from biomass residue: Influence of process conditions to adsorbent properties. *ACS Omega* 6(15): 10224–10233. <https://doi.org/10.1021/acsomega.1c00530>
- Gebrechristos H, Chen W (2018) Utilization of potato peel as eco-friendly products: A review. *Food Sci and Nutr* 6(6): 1352–1356. <https://doi.org/10.1002/fsn3.691>
- Gęca M, Wiśniewska M, Nowicki P (2022) Biochars and activated carbons as adsorbents of inorganic and organic compounds from multicomponent systems – A review. *Adv Colloid Interface Sci* 305: 102–117. <https://doi.org/10.1016/J.CIS.2022.102687>
- Goyi A, Mohammad N, Omer, K (2023) Preparation and characterization of potato peel derived hydrochar and its application for removal of Congo red: A comparative study with potato peel powder. *IJEST* 21(1): 631–642. <https://doi.org/10.1007/S13762-023-04965-Y>
- Haque M, Ibrahim G, Sundarrajan P (2022) Extraction of antioxidants from potato peels and incorporation into value-added products. *AP* 11(1): 389–395. <https://doi.org/10.54085/ap.2022.11.1.44>
- Helal M, Adawy T, Beltagy A, Bedawey A, Youssef S (2020) Evaluation of potato peel extract as a source of antioxidant and antimicrobial substances. *MJFDS* 5(6): 79–90. <https://doi.org/10.21608/mjfds.2020.171473>
- Hoseinzadeh E, Samarghandi M, McKay G, Rahimi N, Jafari J (2014) Removal of acid dyes from aqueous solution using potato peel waste biomass: A kinetic and equilibrium study. *Desalin* 52(27): 4999–5006. <https://doi.org/10.1080/19443994.2013.810355>

- Javed A, Ahmad A, Tahir A, Shabbir U, Nouman M, Hameed A (2019) Potato peel waste—its nutraceutical, industrial and biotechnological applications. *AIMS Agric Food* 4(3): 807–823. <https://doi.org/10.3934/agrfood.2019.3.807>
- John D, Hussin N, Shahibi M, Ahmad M, Hashim H, Ametefe D (2023) A systematic review on the factors governing precision agriculture adoption among small-scale farmers 52(4): 469–485. <https://doi.org/10.1177/00307270231205640>
- Kalio G, Emeya S, Okafor B (2015) Availability and utilization of crop by-products as livestock feeds for small ruminants in khana local government area, rivers state, Nigeria. *Asian j agric* 7(3): 1–9. <https://doi.org/10.9734/ajaees/2015/16203>
- Kayiwa R, Kasedde H, Lubwama M, Kirabira J (2021) The potential for commercial scale production and application of activated carbon from cassava peels in Africa: A review. *Bioresour Technol* 15: 92–106. <https://doi.org/10.1016/j.biteb.2021.100772>
- Kayiwa R, Kasedde H, Lubwama M, Kirabira J (2022) Active pharmaceutical ingredients sequestered from water using novel mesoporous activated carbon optimally prepared from cassava peels. *Water* 14(21): 3371–3391. <https://doi.org/10.3390/W14213371/S1>
- Khanal S, Karimi K, Majumdar S, Kumar V, Verma R, Bhatia S, Kuca K, Esteban J, Kumar D (2023) Sustainable utilization and valorization of potato waste: State of the art, challenges, and perspectives. *Biomass Convers Biorefin* 1: 1–26. <https://doi.org/10.1007/S13399-023-04521-1>
- Kumar V, Othman N, Asharuddin S (2017) Applications of natural coagulants to treat wastewater – a review. *MATEC Web Conf* 103: 6016–6025. <https://doi.org/10.1051/MATECCONF/201710306016>
- Kumar V, Al-Gheethi A, Asharuddin S, Othman N (2021) Potential of cassava peels as a sustainable coagulant aid for institutional wastewater treatment: Characterisation, optimisation and techno-economic analysis. *J Chem Eng* 420: 1–50. <https://doi.org/10.1016/j.cej.2020.127642>
- Lal M, Tiwari R, Kumar R, Naga K, Kumar A, Singh B, Raigond P, Dutt S, Chourasia K, Kumar D, Parmar V, Changan, S (2021) Effect of potato apical leaf curl disease on glycemic index and resistant starch of potato (*Solanum tuberosum L.*) tubers. *Food Chem* 359: 939–948. <https://doi.org/10.1016/j.foodchem.2021.129939>
- Lastochkina O, Baymiev A, Shayahmetova A, Garshina D, Koryakov I, Shpirnaya I, Pusenkova L, Mardanshin I, Kasnak C, Palamutoglu, R (2020) Effects of endophytic bacillus subtilis and salicylic acid on postharvest diseases (*Phytophthora infestans*, *fusarium oxysporum*) development in stored potato tubers. *Plants* 9(1): 76–98. <https://doi.org/10.3390/plants9010076>
- Li D, Zhao X, Liu Z, Liu H, Fan B, Yang B, Zheng X, Li W, Zou H (2021) Synergetic anticorrosion mechanism of main constituents in Chinese yam peel for copper in artificial seawater. *ACS Omega* 6(44): 29965–29981. <https://doi.org/10.1021/acsomega.1c04500>
- Liang S, Han Y, Wei L, McDonald A (2015) Production and characterization of bio-oil and bio-char from pyrolysis of potato peel wastes. *Biomass Convers Biorefin* 5(3): 237–246. <https://doi.org/10.1007/s13399-014-0130-x/metrics>
- Mahapatra D, Satapathy K, Panda B (2022) Biofertilizers and nanofertilizers for sustainable agriculture: Phyco-prospects and challenges. *Sci Total Environ* 803: 376–393. <https://doi.org/10.1016/j.scitotenv.2021.149990>

- Miller K, Reichert C, Schmid M, Loeffler M (2022) Physical, chemical and biochemical modification approaches of potato (peel) constituents for bio-based food packaging concepts: A review. *Foods* 11(18): 2927–2965. <https://doi.org/10.3390/foods11182927>
- Min S, Ezati P, Rhim, J (2022) Gelatin-based packaging material incorporated with potato skins carbon dots as functional filler. *Industrial Crops and Products* 181: 128–141. <https://doi.org/10.1016/j.indcrop.2022.114820>
- Minisha S, Vedhi C, Rajakani P, Pertiwi D, Yanti N, Taslim R (2022) High potential of yellow potato (*Solanum Tuberosum L.*) peel waste as porous carbon source for supercapacitor electrodes. *J Phys Conf Ser* 2193(1): 1–11. <https://doi.org/10.1088/1742-6596/2193/1/012019>
- Moretti B, Bertora C, Grignani C, Lerda C, Celi L, Sacco D (2020) Conversion from mineral fertilisation to MSW compost use: Nitrogen fertiliser value in continuous maize and test on crop rotation. *Sci Total Environ* 705: 27–58. <https://doi.org/10.1016/j.scitotenv.2019.135308>
- Mourão M, Xavier L, Urbatzka R, Figueiroa L, Costa C, Dias C, Schneider M, Vasconcelos V, Santos A (2021) Characterization and biotechnological potential of intracellular polyhydroxybutyrate by *stigeoclonium* sp. b23 using cassava peel as carbon source. *Polym* 13(5): 687–708. <https://doi.org/10.3390/polym13050687>
- Muhammad F, Abdul R, Michele P (2019) Sustainable agriculture and food security. In *innovations in sustainable agriculture*. Springer International Publishing, New York, pp. 3–24. https://doi.org/10.1007/978-3-030-23169-9_3
- Mulia D, Raicha R, Lunggani C, Erina S, Wuliandari J, Purbomartono C, Isnansetyo A (2023) Antibacterial activity of ethanol extract of banana, cassava, and pineapple peels against a fish pathogen *Aeromonas hydrophila*. *Biodivers J* 24(1): 481–485. <https://doi.org/10.13057/biodiv/d240155>
- Nahas S, Salman H, Ahmed W, Salman H, Seleeme W (2019) Aluminum building scrap wire, take-out food container, potato peels and bagasse as valueless waste materials for nitrate removal from water supplies. *Chem Afr* 2(1): 143–162. <https://doi.org/10.1007/s42250-018-00032-z>
- Nair L, Agrawal K, Verma P (2022) An overview of sustainable approaches for bioenergy production from agro-industrial wastes. *J Nexus* 6: 1–25. <https://doi.org/10.1016/J.NEXUS.2022.100086>
- Nathan R, Martin C, Barr D, Rosengren R (2021) Simultaneous removal of heavy metals from drinking water by banana, orange and potato peel beads: A study of biosorption kinetics. *Appl Water Sci* 11(7): 1–15. <https://doi.org/10.1007/S13201-021-01457-7>
- Neela S, Fanta S (2019) Review on nutritional composition of orange-fleshed sweet potato and its role in management of vitamin A deficiency. *Food Sci Nutr* 7(6): 1920–1945. <https://doi.org/10.1002/FSN3.1063>
- Nemadziva B, Ngubane S, Matiza Ruzengwe F, Kasumbwe K, Kudanga T (2022) Potato peels as feedstock for laccase-catalysed synthesis of *phellinsin* A. *Biomass Convers Biorefin* 13(15): 13871–13882. <https://doi.org/10.1007/S13399-021-02251-W>
- Odeyemi S, Iwuzor K, Emenike E, Odeyemi O, Adeniyi A (2023) Valorization of waste cassava peel into biochar: An alternative to electrically-powered process. *Totert* 6: 1–10. <https://doi.org/10.1016/j.totert.2023.100029>
- Oghenejoboh K, Orugba H, Oghenejoboh M, Agarry S (2021) Value added cassava waste management and environmental sustainability in Nigeria: A review. *Environ* 4: 1–14. <https://doi.org/10.1016/j.envc.2021.100127>
- Ojuederie O, Olanrewaju O, Babalola O (2019) Plant growth promoting rhizobacterial mitigation of drought stress in crop plants: implications for sustainable agriculture. *Agron* 9(11): 712–741. <https://doi.org/10.3390/agronomy9110712>

- Okekunle P, Itabiyi O, Adetola S, Alayande I, Ogundiran H, Odeh K (2016) Biofuel production by pyrolysis of cassava peel in a fixed bed reactor. *Int J Energy Clean Environ* 17(1): 57–65. <https://doi.org/10.1615/interjenercleanenv.2017018176>
- Okike I, Samireddypalle A, Kaptoge L, Fauquet C, Atehnkeng J, Bandyopadhyay R, Kulakow P, Duncan A, Alabi T, Blummel M (2015) Technical innovations for small-scale producers and households to process wet cassava peels into high quality animal feed ingredients and aflasafe substrate. *Food Chain* 5(1): 71–90. <https://doi.org/10.3362/2046-1887.2015.005>
- Okike I, Seerp W, Anandan S, Diego N, Peter K (2022) Turning waste to wealth: Harnessing the potential of cassava peels for nutritious animal feed. In *root, tuber and banana food system innovations*. Springer International Publishing, New York, pp. 173–206. <https://doi.org/10.1007/978-3-030-92022-7>
- Oladele I, Ibrahim I, Adediran A, Akinwekomi A, Adetula Y, Olayanju T (2020) Modified palm kernel shell fiber/particulate cassava peel hybrid reinforced epoxy composites. *Results Mater* 5: 1–7. <https://doi.org/10.1016/j.rinma.2019.100053>
- Oloruntola O (2020) Effect of dietary cassava peel meal supplemented with methionine and multienzyme on hemobiochemical indices, digestibility, and antioxidants in rabbits. *JOBAZ* 81(1):1–11. <https://doi.org/10.1186/s41936-020-00170-2>
- Omole A, Okpeze C, Fayenuwo J, Olorungbohunmi T (2013) Effects of partial replacement of maize with yam peel (*Discorea rotundata*) in diet of juvenile snails (*Archachatina marginata*). *Afr J Agric Res* 8(16): 1361–1364. <https://doi.org/10.5897/ajar11.1490>
- Onu C, Nweke C, Nwabanne J (2022) Modeling of thermo-chemical pretreatment of yam peel substrate for biogas energy production: RSM, ANN, and ANFIS comparative approach. *Appl Surf Sci* 11: 1–13. <https://doi.org/10.1016/j.apsadv.2022.100299>
- Osemwengie S, Osagie E, Onwukwe B (2020) Optimization of bioethanol production from cassava peels. *J Appl Sci Environ* 24(12): 2077–2083. <https://doi.org/10.4314/jasem.v24i12.11>
- Osman A, Blewitt J, Abu-Dahrieh J, Farrell C, Al-Muhtaseb A, Harrison J, Rooney D (2019) Production and characterisation of activated carbon and carbon nanotubes from potato peel waste and their application in heavy metal removal. *Environ Sci Pollut Res* 26(36): 37228–37241. <https://doi.org/10.1007/S11356-019-06594>
- Ospino J, Barraza J, Cervera S, Escobar E, Ceballos O, Ospino J, Barraza J, Cervera S (2022) Activated carbon from cassava peel: A promising electrode material for supercapacitors. *Rev Fac Ing Univ Ant* 102: 88–95. <https://doi.org/10.17533/udea.redin.20200803>
- Page M, McKenzie J, Bossuyt P, Boutron I, Hoffmann T, Mulrow C, Shamseer L, Tetzlaff J, Akl E, Brennan S, Chou R, Glanville J, Grimshaw J, Hróbjartsson A, Lalu M, Li T, Loder E, Wilson E, McDonald S, Moher D (2021) The Prisma 2020 statement: An updated guideline for reporting systematic reviews. *Int Surg J* 88: 1–9. <https://doi.org/10.1016/j.ijju.2021.105906>
- Pang W, Hou D, Chen J, Nowar E, Li Z, Hu R, Tomberlin J, Yu Z, Li Q, Wang S (2020) Reducing greenhouse gas emissions and enhancing carbon and nitrogen conversion in food wastes by the black soldier fly. *J Environ Manag* 260: 1–8. <https://doi.org/10.1016/j.jenvman.2020.110066>
- Papathoti N, Laemchiab K, Megavath V, Keshav P, Numparditsub P, Thanh T, Buensanteai N (2021) Augmented ethanol production from alkali-assisted hydrothermal pretreated cassava peel waste. *Energy Sources A: Recovery Util Environ* 20: 1–11. <https://doi.org/10.1080/15567036.2021.1928338>

- Phestrungnapha A, Wiengnak N, Maikrang K (2023) Removal of free fatty acid from waste cooking oil using an adsorbent derived from cassava peels. *Korean J Chem Eng* 40(9): 2253–2262. <https://doi.org/10.1007/S11814-023-1413-3>
- Pulungan M, Kapita R, Dewi I (2020) Optimisation on the production of biodegradable plastic from starch and cassava peel flour using response surface methodology. *IOP Conf Ser Earth Environ Sci* 475(1): 1–9. <https://doi.org/10.1088/1755-1315/475/1/012019>
- Rahman M, Mat K, Ishigaki G, Akashi R (2021) A review of okara (soybean curd residue) utilization as animal feed: Nutritive value and animal performance aspects. *Anim Sci J* 92(1): 1–8. <https://doi.org/10.1111/asj.13594>
- Rahmani M, Azadbakht M, Dastar B, Esmaeilzadeh E (2022) Production of animal feed from food waste or corn? Analyses of energy and exergy. *Bioresour Technol Rep* 20: 101213–101221. <https://doi.org/10.1016/j.biteb.2022.101213>
- Redlingshöfer B, Barles S, Weisz H (2020) Are waste hierarchies effective in reducing environmental impacts from food waste? A systematic review for OECD countries. *Resour Conserv Recycl* 156: 1–17. <https://doi.org/10.1016/j.resconrec.2020.104723>
- Rengsutthi K, Charoenrein S (2011) Physico-chemical properties of jackfruit seed starch (*Artocarpus heterophyllus*) and its application as a thickener and stabilizer in chilli sauce. *LWT - Food Sci* 44(5): 1309–1313. <https://doi.org/10.1016/j.lwt.2010.12.019>
- Rinaldo D (2020) Carbohydrate and bioactive compounds composition of starchy tropical fruits and tubers, in relation to pre and postharvest conditions: A review. *J Food Sci* 85(2): 249–259. <https://doi.org/10.1111/1750-3841.15002>
- Rodríguez B, Gullón B, Yáñez R (2021) Identification and recovery of valuable bioactive compounds from potato peels: A comprehensive review. *Antiox* 10(10): 1630–1641. <https://doi.org/10.3390/antiox10101630>
- Ruqayyah T, Jamal P, Alam M, Mirghani M, Jaswir I, Ramli N (2014) Application of response surface methodology for protein enrichment of cassava peel as animal feed by the white-rot fungus *Panus tigrinus* M609RQY. *Food Hydrocol* 42(P2): 298–303. <https://doi.org/10.1016/j.foodhyd.2014.04.027>
- Saeed A, Shabbir A, Khan A (2022) Stabilization of sunflower oil by using potato peel extract as a natural antioxidant. *Biomass Conv Bioref* 14: 5275–5284. <https://doi.org/10.1007/S13399-022-02696-7>
- Samotya U (2019) Potato peel as a sustainable resource of natural antioxidants for the food industry. *Potato Res* 62(4): 435–451. <https://doi.org/10.1007/S11540-019-9419-2>
- Sampaio S, Petropoulos S, Alexopoulos A, Heleno S, Buelga C, Barros L, Ferreira I (2020) Potato peels as sources of functional compounds for the food industry: A review. *Trends Food Sci Technol* 103: 118–129. <https://doi.org/10.1016/j.tifs.2020.07.015>
- Sarwono R (2023) Approaching zero waste management of municipal solid waste integrated with agricultural, poultry, ruminants, worm and maggot farming. *Int J Eng Sci* 12(7): 6–10. <https://doi.org/10.9790/1813-12070610>
- Scott G (2021) A review of root, tuber and banana crops in developing countries: Past, present and future. *Int. J Food Sci Technol* 56(3): 1093–1114. <https://doi.org/10.1111/ijfs.14778>
- Shalini S, Raghavan V (2021) Biochar from biomass waste as a renewable carbon material for climate change mitigation in reducing greenhouse gas emissions - a review. *Biomass Convers Biorefin* 1(11): 2247–2267. <https://doi.org/10.1007/s13399-020-00604-5>

- Shao Y, Zheng C, Liu K, Xiong J, Wang X, Han M, Li L, Shi Y, Lu J, Yi J (2022) Extraction optimization, purification, and biological properties of polysaccharide from Chinese yam peel. *J Food Biochem* 46(12): 147–152. <https://doi.org/10.1111/jfbc.14490>
- Singh L, Kaur S, Aggarwal P, Kaur N (2023) Characterisation of industrial potato waste for suitability in food applications. *Int J Food Sci Technol* 58(5): 2686–2694. <https://doi.org/10.1111/ijfs.16023>
- Singh M, Rano S, Roy S, Mukherjee P, Dalui S, Gupta G, Kumar S, Mondal M (2022) Characterization of organophosphate pesticide sorption of potato peel biochar as low-cost adsorbent for chlorpyrifos removal. *Chemosphere* 297: 43–54. <https://doi.org/10.1016/j.chemosphere.2022.134112>
- Sugumaran V, Vimal K, Kapur G, Narula A (2015) Preparation and morphological, thermal, and physicochemical properties of polypropylene-potato peel biocomposites. *J Appl Polym Sci* 132(34): 1–12. <https://doi.org/10.1002/APP.42445>
- Sui B, Meng H, Shen Y, Ding J, Wang J (2018) Utilization of livestock manure in Denmark and its inspiration for planting-breeding combined circular agricultural development in China. *Nongye Gongcheng Xuebao Trans Chin Soc Agric Eng* 34(12): 1–7. <https://doi.org/10.11975/J.ISSN.1002-6819.2018.12.001>
- Surai P (2014) Polyphenol compounds in the chicken/animal diet: From the past to the future. *J Anim Physiol Anim Nutr* 98(1): 19–31. <https://doi.org/10.1111/jpn.12070>
- Tadele Z (2019) Orphan crops: Their importance and the urgency of improvement. *Planta* 250(3): 677–694. <https://doi.org/10.1007/S00425-019-03210-6>
- Thuppahige V, Moghaddam L, Welsh Z, Wang T, Karim A (2023) Investigation of critical properties of Cassava (*Manihot esculenta*) peel and bagasse as starch-rich fibrous agro-industrial wastes for biodegradable food packaging. *Food Chem* 422: 1–12. <https://doi.org/10.1016/j.foodchem.2023.136200>
- Triwidatin Y, Djuanda U (2023) Creative products from cassava peel waste to be delicious and nutritious snacks, desa sukamanah, megamendung district, bogor regency. *JPMB* 2(6): 455–462. <https://doi.org/10.55927/jpmb.v2i6.4469>
- Vaitkevičienė N (2019) A comparative study on proximate and mineral composition of coloured potato peel and flesh. *J Sci Food Agric* 99(14): 6227–6233. <https://doi.org/10.1002/jsfa.9895>
- Velasco J, Sánchez J, Felices B, Sánchez I (2022) Circular economy in agriculture: An analysis of the state of research based on the life cycle. *Sustain Prod Consum* 34: 257–270. <https://doi.org/10.1016/j.spc.2022.09.017>
- Versino F, López O, García M (2015) Sustainable use of cassava (*Manihot esculenta*) roots as raw material for biocomposites development. *Ind Crops Prod* 65: 79–89. <https://doi.org/10.1016/j.indcrop.2014.11.054>
- Wanapat M, Kang S, Polyorach S (2013) Development of feeding systems and strategies of supplementation to enhance rumen fermentation and ruminant production in the tropics. *J Animal Sci Biotechnol* 4(32): 1–11. <https://doi.org/10.1186/2049-1891-4-32>
- Wang H, Liu S, Zhou X, Yang X, Gao Q, Tanokura M, Xue Y (2020a) Treatment with hydrogen peroxide improves the physicochemical properties of dietary fibres from Chinese yam peel. *Int J Food Sci Technol* 55(3): 1289–1297. <https://doi.org/10.1111/ijfs.14405>
- Wang S, Lin A, Han Q, Xu Q (2020b) Evaluation of direct ultrasound-assisted extraction of phenolic compounds from potato peels. *Processes* 8(12): 1–14. <https://doi.org/10.3390/pr8121665>
- Wolfswinkel J, Furtmueller E, Wilderom C (2013) Using grounded theory as a method for rigorously reviewing literature. *Eur J Inf Syst* 22(1): 45–55. <https://doi.org/10.1057/ejis.2011.51>

- Wu W, Hung W, Lo K, Chen Y, Wan H, Cheng K (2016) Bioethanol production from taro waste using thermo-tolerant yeast *Kluyveromyces marxianus* K21. *Bioresour Technol* 201: 27–32. <https://doi.org/10.1016/j.biortech.2015.11.015>
- Xie F, Zhang W, Lan X, Gong S, Wu J, Wang Z (2018) Effects of high hydrostatic pressure and high-pressure homogenization processing on characteristics of potato peel waste pectin. *Carbohydr Polym* 196: 474–482. <https://doi.org/10.1016/J.carbpol.2018.05.061>
- Yafetto L, Odamtten G, Kwagyan M (2023) Valorization of agro-industrial wastes into animal feed through microbial fermentation: A review of the global and Ghanaian case. *Heliyon* 9(4): 1–14. <https://doi.org/10.1016/j.heliyon.2023.e14814>
- Zierer W, Rüscher D, Sonnewald U, Sonnewald S (2021) Annual review of plant biology tuber and tuberous root development. *Annu Rev Plant Biol* 72: 551–580. <https://doi.org/10.1146/annurev-arplant-080720>