

DOI: [10.57647/pibm-2025-17593](https://doi.org/10.57647/pibm-2025-17593)

SUPPORTING INFORMATION

Encapsulation of IR783 in UiO-66 MOFs for Improved Photodynamic Efficacy Against Breast Cancer Cells

Sinem Şahinoğlu and Gülşah Şanlı-Mohamed*

Department of Chemistry, İzmir Institute of Technology, 35430, İzmir, Türkiye

*Corresponding Author

Prof. Dr. Gülşah Şanlı Mohamed

İzmir Institute of Technology, Science Faculty, Department of Chemistry
İzmir, Türkiye

Phone: +90 2327507515; Fax: +90 2327507509

E-mail: gulsahsanli@iyte.edu.tr, gulsahsanli@gmail.com




Figure S1. Cytotoxicity profiles of IR783, UiO-66 and IR783@UiO-66 in MCF-10A cell line at 12 and 48 h.




Figure S2. Phototoxicity profiles of IR783, UiO-66 and IR783@UiO-66 in MCF-10A cell line after 24 h of irradiation under LED (18, 36 and 72 J/cm²).




Figure S3. Cytotoxicity profiles of IR783, UiO-66 and IR783@UiO-66 in MDA-MB-231 cell line at 12 and 48 h.




Figure S4. Phototoxicity profiles of IR783, UiO-66 and IR783@UiO-66 in MDA-MB-231 cell line after 24 h of irradiation under LED (18, 36 and 72 J/cm²).




Figure S5. Cytotoxicity profiles of IR783, UiO-66 and IR783@UiO-66 in MCF-7 cell line at 12 and 48 h.




Figure S6. Phototoxicity profiles of IR783, UiO-66 and IR783@UiO-66 in MCF-7 cell line after 24 h of irradiation under LED (18, 36 and 72 J/cm²).